arXiv:1304.1076v1 [math.DG] 3 Apr 2013

Pure spinors, intrinsic torsion and curvature in odd dimensions

Arman Taghavi-Chabert
Masaryk University, Faculty of Science, Department of Mathematics and Statistics,
Kotlářská 2, 611 37 Brno, Czech Republic

Abstract

We develop a spinor calculus for a $(2m+1)$-dimensional complex Riemannian manifold $(\mathcal{M}, g)$ equipped with a preferred holomorphic projective pure spinor field $[\xi]$. Such a spinor defines a holomorphic distribution $\mathcal{N}$ of m-planes, totally null with respect to g. The stabiliser of $[\xi]$ is a parabolic Lie subgroup P of $\mathrm{SO}(2m+1, \mathbb{C})$, and using its algebraic properties, we give P-invariant classifications of the curvature tensors of the Levi-Civita connection, which, in the case of the Weyl tensor, generalises the Petrov-Penrose classification to odd dimensions. From a spinorial point of view, this generalises the notion of principal spinors.

We also classify the intrinsic torsion of $[\xi]$ in terms of P-irreducibles, which gives an algebraic measure of the failure of $[\xi]$ to be parallel with respect to the Levi-Civita connection. In particular, we interpret the integrability properties of the associated null m-plane distribution $\mathcal{N}$ and its orthogonal complement $\mathcal{N}^\perp$ in terms of the algebraic types of the intrinsic torsion of $[\xi]$. The conformal invariance of these classes are studied and we give curvature conditions for a number of cases.

We then investigate the geometric properties of a number of spinorial differential equations. Notably, we relate the integrability properties of a null m-plane distribution to the existence of solutions of odd-dimensional versions of the zero-rest-mass field equation. We also give necessary and sufficient conditions for a pure conformal Killing spinor to be foliating. This leads in particular to a conjecture refining the author's odd-dimensional generalisation of the Goldberg-Sachs theorem.

This work can also be applied to a smooth real pseudo-Riemannian manifold of split signature equipped with a preferred projective real pure spinor field.

1 Introduction and motivation

The present article is the 'odd-dimensional counterpart' of the author's work presented in [TC12b], which focused on the geometric properties of projective pure spinor fields in even dimensions. Both this article and [TC12b] work share the same motivations and goals, and the reader should refer to the latter work for further details.

The introduction of spinors in [Wit59] and [Pen60] to formulate Petrov's classification of the Weyl tensor in Lorentzian signature [Pet00] marks an important step in the understanding of the complex geometric properties of spacetimes. This approach impinges on the fact that in four dimensions, Lorentzian signature, the self-dual part and anti-self-dual part of the Weyl tensor are complex conjugate of each other. Thus, the Petrov classification of the Weyl tensor can be reduced to the classification of its complex self-dual part. Algebraically, this can be represented by an element of the fourth-symmetric-power $\odot^4 \mathfrak{S}^+$ of the two-dimensional complex representation $\mathfrak{S}^+$ of positive spinors. This element can be thought of as the components of a homogeneous polynomial of degree 4, and the multiplicities of its roots gives an invariant way of classifying the Weyl tensor. A root of this polynomial is essentially a positive spinor $[\xi]$ up to scale, i.e. a point in $\mathbb{CP}^1$, and such a spinor is known as a *gravitational principal spinor*. When the Weyl tensor (at a point) admits a repeated root, it is then said to be *algebraically special*. This is known as the Petrov-Penrose type $\{2,1,1\}$ or type II.

In general relativity, algebraically special solutions to Einstein's field equations play an important geometric rôle: by the Goldberg-Sachs theorem [GS09], an Einstein spacetime is algebraically special if and only if it admits a *shearfree congruence of null geodesics (SCNG)*. It is, however, more convenient to understand a SCNG from the perspective of spinors. A projective chiral spinor field defines a distribution $\mathcal{N}$, say, of complex 2-planes, totally null with respect to the complexified metric. The integrability of $\mathcal{N}$ is then *equivalent* to the existence of a SNCG, which can be seen to arise from the intersection of $\mathcal{N}$ and its complex conjugate $\bar{\mathcal{N}}$.

The advantage of such an interpretation is that it easily adapts to other metric signatures. If our underlying manifold is not Lorentzian, we then have two independent Petrov-Penrose classifications corresponding to the self-dual part and anti-self-part of the Weyl tensor. The Goldberg-Sachs theorem still holds, but each signature yields a different geometric interpretation of what a foliating spinor is. For instance, in Euclidean signature, a totally null complex 2-plane distribution and its complex conjugate are simply the $+\mathrm{i}$-eigenspace and $-\mathrm{i}$-eigenspace of an almost Hermitian structure respectively so that algebraically special Einstein Riemannian manifolds locally admit complex structures [AG97, GHN10]. On the other hand, if the underlying manifold is a *complex Riemannian manifold*, i.e. a complex manifold endowed with a holomorphic non-degenerate symmetric bilinear form, on which no particular reality structure is assumed, one only has a *single* spinor field defining a totally null 2-plane distribution. This is in a sense the 'barest' and most transparent setting, and will be the focus of this paper.

One way to generalise this to complex Riemannian manifolds of dimension $2m+1$ is to consider a *γ-plane distribution*, i.e. a holomorphic distribution $\mathcal{N}$ of m-planes that are totally null with respect to the metric, or equivalently, a holomorphic projective spinor field $[\xi]$, say, annihilated by this distribution – such a spinor is said to be *pure* [Car81]. The existence of such a spinor field reduces the structure group of the frame bundle to its stabiliser P, say, a subgroup of the complex special orthogonal group $\mathrm{SO}(2m+1,\mathbb{C})$. The subgroup P is a well-known example of a *parabolic Lie subgroup* [BE89, FH91, ČS09b]. This naturally leads to an algebraic classification of the Weyl tensor, and other irreducible curvature tensors, invariant under P: the space of tensors with Weyl symmetries $\mathfrak{C}$ admits a filtration $\mathfrak{C}^4 \subset \mathfrak{C}^3 \subset \ldots \mathfrak{C}^{-3} \subset \mathfrak{C}^{-4} := \mathfrak{C}$ of P-modules $\mathfrak{C}^i$. One can then go a step further by decomposing each P-module $\mathfrak{C}^i/\mathfrak{C}^{i+1}$ into irreducibles, each being isomorphic to an irreducible $\mathrm{GL}(m,\mathbb{C})$-module, the reductive part of P.

In fact, a partial higher-dimensional generalisation of the Goldberg-Sachs theorem was presented in [TC12a]: therein a notion of algebraic special complex Riemannian manifolds based on the filtration $\{\mathfrak{C}^i\}$ is defined, which under further genericity assumptions on the Weyl tensor and degeneracy assumptions on the Cotton-York tensor, guarantees the integrability of both $\mathcal{N}$ *and* its orthogonal complement.[1] Such a result however remains coarse on account of the genericity assumption, and as we shall see in section 5, a pure conformal Killing spinor may not be foliating, yet its integrability condition implies the algebraic speciality of the Weyl tensor in the sense of [TC12a]. One must therefore seek weaker differential conditions on the distribution $\mathcal{N}$ (or its associated projective spinor field $[\xi]$), and for this purpose, we turn our attention to the *intrinsic torsion* of the P-structure of the frame bundle [Sal89]. This is a tensorial quantity identified with the Levi-Civita covariant derivative $\nabla[\xi]$ of $[\xi]$, and lies in the P-module $\mathfrak{W} := \mathfrak{V} \otimes (\mathfrak{g}/\mathfrak{p})$ – here, $\mathfrak{g} := \mathfrak{so}(2m+1,\mathbb{C})$ is the Lie algebra of $\mathrm{SO}(2m+1,\mathbb{C})$, and $\mathfrak{V}$ its standard representation. The decomposition of $\mathfrak{W}$ into irreducibles P-modules then provides a natural scheme under which differential conditions on the distribution $\mathcal{N}$ and its orthogonal complement can be formulated.

The aim of this article is to refine the classification of [TC12a] and to extend the work of [TC12b] to odd dimensions. In particular, given a projective pure spinor $[\xi]$, we shall develop a spinor calculus in order to

- formulate a spinorial classification of irreducible curvature tensors such as the Weyl tensor generalising the Petrov-Penrose classification;
- characterise some of the geometric properties of the γ-plane distribution of $[\xi]$ in terms of its intrinsic torsion, i.e. algebraic conditions on $\nabla[\xi]$;
- and apply this machinery to a number of spinorial differential equations such as the zero-rest-mass field equation and the twistor equation.

---

[1] This was motivated by the Lorentzian version given in [TC11], where it was shown that such conditions naturally lead to the existence of a congruence of null *geodesics*. Also, this property was shown to feature in a number of higher-dimensional solutions of Einstein's field equations - see also [MT10].

The differential properties of pure spinor fields appear to have first been investigated in six and higher even dimensions in [Hug95, Jef95, HM88]. The reference [Jef95] in fact contains a prototypical spinorial classification of the Weyl tensor in six dimensions. In five dimensions, the author of [DS05] put forward a spinorial classification of the Weyl tensor distinct from the one presented in the present article. It stems from the remark that in five dimensions, the Weyl tensor is a totally symmetric 4-spinor, from which one can construct a homogeneous polynomial of degree 4 in $\mathbb{CP}^3$. This was thoroughly investigated in [God10] in Lorentzian signature. A five-dimensional spinor calculus was also introduced in [GPGLMG09].

The classifications of the intrinsic torsion and the curvature of almost Hermian manifolds were given in [GH80] and [TV81, FFS94] respectively. The application of this work and [TC12b] to the study of Lorentzian geometry will be presented in [TC], and can be thought of as a refinement of the higher-dimensional Petrov-Weyl classification expounded in [CMPP04, MCPP05].

The structure of this article follows that of [TC12b]. In section 2, we recall some background on Clifford algebras and its spinor representation. Fixing a preferred pure spinor, we build a spinor calculus emphasising its relation to the representation theory of its stabiliser, a parabolic Lie subalgebra $\mathfrak{p}$ of $\mathfrak{g} = \mathfrak{so}(2m+1, \mathbb{C})$. This leads, in section 3, to the $\mathfrak{p}$-invariant classifications of the tracefree Ricci tensor, the Cotton-York tensor and the Weyl tensor leading to Propositions 3.1, 3.2 and 3.3 respectively, which include 'Penrose diagrams' for these tensors.

In section 4, we classify the module $\mathfrak{W} = \mathfrak{V} \otimes (\mathfrak{g}/\mathfrak{p})$ in terms of irreducible $\mathfrak{p}$-modules as given by Proposition 4.2. This paves the way for the classification of the intrinsic torsion of a projective pure spinor field given in the next section.

Section 5 focuses on the geometric applications of the algebraic classifications of the previous sections. Building on section 4, we give algebraic characterisations of the intrinsic torsion of a projective pure spinor field $[\xi]$ in Proposition 5.4. The integrability properties of the null m-plane distribution of $[\xi]$ and its orthogonal complement are interpreted as conditions on the intrinsic torsion of $[\xi]$ in Proposition 5.5. The conformal invariance of the intrinsic torsion is examined in Proposition 5.8, which also includes an odd-dimensional complex Riemannian of the 'conformal to Kähler condition'.

We end the section by investigating the relation between differential equations on pure spinor fields and the differential properties of their associated γ-plane distributions. More specifically, Theorems 5.17, 5.18 and 5.19 give odd-dimensional versions of the Robinson theorem for three distinct types of zero-rest-mass fields. Further, Proposition 5.21 and 5.24 investigate the differential properties of the γ-plane distribution associated to a pure conformal Killing spinor. Finally, we give a conjecture 5.27 of the Goldberg-Sachs theorem, generalising that of [TC12a].

We specialise to the special three- and five-dimensional cases in appendix B, where a number of results in the main text simplify. In appendix C, we collect some useful formulae regarding conformal geometry. Finally, we have collected tables of irreducibles in appendix D.

2 Algebraic background

In this section, we review the background on Clifford algebras, its spinor representations, and the algebraic properties of pure spinors. The approach will be to synthesize aspects of the theory of spinors given notably in [Car81, PR86, BT89, HS95, Kop97] with the representation theory of parabolic Lie subalgebras of [BE89, ČS09b]. We shall adopt the abstract index notation of [PR84, PR86] throughout, in particular, that of Clifford algebras and spinors.

2.1 Clifford algebras and spinor representations

Let $\mathfrak{V}$ be a n-dimensional complex vector space. Elements of $\mathfrak{V}$, respectively, of its dual $\mathfrak{V}^*$, will carry upstairs, respectively, downstairs, lower-case Roman indices, e.g. V^a and α_a respectively. This notation extends to tensor products of $\mathfrak{V}$ and $\mathfrak{V}^*$, i.e. we write $T_{ab}{}^c{}_d$ for an element of $\otimes^2\mathfrak{V}^* \otimes \mathfrak{V} \otimes \mathfrak{V}^*$. We equip $\mathfrak{V}$ with a non-degenerate symmetric bilinear form $g_{ab} = g_{(ab)} \in \odot^2\mathfrak{V}^*$. Here, as elsewhere, symmetrisation is denoted by round brackets, while skew-symmetrisation by squared brackets, e.g. $\alpha_{abc} = \alpha_{[abc]} \in \wedge^3\mathfrak{V}^*$. The inner product g_{ab} together with its inverse g^{ab} establishes an isomorphism between $\mathfrak{V}$ and $\mathfrak{V}^*$, so that one will lower or raise the indices of tensorial quantities as needed.

The Clifford algebra $\mathcal{C}\ell(\mathfrak{V},g)$ of $(\mathfrak{V},g)$ is defined as the quotient algebra $\bigotimes^\bullet \mathfrak{V}/\mathfrak{I}$ where $\mathfrak{I}$ is the ideal generated by elements of the form $v \otimes v + g(v,v)$ where $v \in \mathfrak{V}$. This means that $\mathcal{C}\ell(\mathfrak{V},g)$ and the exterior algebra $\wedge^\bullet \mathfrak{V}$ are isomorphic as vector spaces, the wedge product being now replaced with the Clifford product $\cdot : \mathcal{C}\ell(\mathfrak{V},g) \times \mathcal{C}\ell(\mathfrak{V},g) \to \mathcal{C}\ell(\mathfrak{V},g)$ defined by

$$v \cdot w := v \wedge w - g(v,w) \,,$$

for $v, w \in \mathfrak{V}$ viewed as elements of $\mathcal{C}\ell(\mathfrak{V},g)$.

Henceforth, we assume $n = 2m+1$, in which case the Clifford algebra can be seen to be isomorphic to a direct sum of two copies of the algebra of $2^m \times 2^m$-matrices over $\mathbb{C}$. Each copy acts on a 2^m-dimensional complex vector space, which is known as the *spinor space of* $(\mathfrak{V},g)$. These spinor spaces are inequivalent irreducible representations of the double cover $\mathrm{Spin}(2m+1,\mathbb{C})$ of $\mathrm{SO}(2m+1,\mathbb{C})$. From now on, we shall work with one of these spinor representations, which we shall denote $\mathfrak{S}$.

Remark 2.1 One can give an explicit construction of the spinor representation $\mathfrak{S}$ in terms of forms as follows. Choose a totally null m-dimensional subspace $\mathfrak{N}$ of $\mathfrak{V}$, and fix a dual $\mathfrak{N}^*$ of $\mathfrak{N}$ so that $\mathfrak{V} \cong \mathfrak{N} \oplus \mathfrak{N}^* \oplus \mathfrak{U}$ where $\mathfrak{U}$ is the (one-dimensional) complement of $\mathfrak{N} \oplus \mathfrak{N}^*$ in $\mathfrak{V}$. Now, for any $(v,w,u) \in \mathfrak{V}$, the action of the Clifford algebra on $\mathfrak{S} \cong \wedge^\bullet \mathfrak{N}$ is given by

$$(v,w,u) \cdot \xi = v \wedge \xi - w \lrcorner \xi \pm \epsilon u \xi \,, \tag{2.1}$$

where

$$\epsilon = \begin{cases} +1 \,, & \text{if } \xi \in \wedge^m \mathfrak{N} \oplus \wedge^{m-2} \mathfrak{N} \oplus \wedge^{m-4} \mathfrak{N} \oplus \ldots \,, \\ -1 \,, & \text{if } \xi \in \wedge^{m-1} \mathfrak{N} \oplus \wedge^{m-3} \mathfrak{N} \oplus \wedge^{m-5} \mathfrak{N} \oplus \ldots \,. \end{cases}$$

The choice of sign in (2.1) reflects the choice of inequivalent spinor representations.

Elements of $\mathfrak{S}$ will carry upstairs upper-case Roman indices, e.g. ξ^A, and similarly for elements of the dual spinor space $\mathfrak{S}^*$, with downstairs indices, e.g. η_A. From the definition of the Clifford algebra, one can show that $\mathcal{C}\ell(\mathfrak{V},g)$ is generated by the *(Van der Waerden)* γ*-matrices* ${\gamma_{aA}}^B$ which satisfy the Clifford property

$$\gamma_{(a\,A}{}^C \gamma_{b)C}{}^B = -g_{ab} \delta_A^B \,. \tag{2.2}$$

Thus, only skew-symmetrised products of γ-matrices count, and we shall make use of the notational short hand

$$\gamma_{a_1 a_2 \ldots a_p A}{}^B := \gamma_{[a_1\,A}{}^{C_1} \gamma_{a_2 C_1}{}^{C_2} \ldots \gamma_{a_p] C_p}{}^B \,. \tag{2.3}$$

These matrices give us an explicit realisation of the isomorphism $\mathcal{C}\ell(\mathfrak{V},g) \cong \bigwedge^\bullet \mathfrak{V}$ as vector spaces.

The spinor space $\mathfrak{S}$ and its dual $\mathfrak{S}^*$ are equipped with non-degenerate bilinear forms, with which one can in effect raise or lower the indices. In particular, the γ-matrices (2.3) can now be viewed as bilinear maps from $\mathfrak{S} \times \mathfrak{S}$ to $\wedge^p \mathfrak{V}$ some p. Depending on the values of m and k, these bilinear forms can be either symmetric or skewsymmetric. Of interest is the case where $p = m$: the γ-matrices $\gamma_{a_1 a_2 \ldots a_m}{}^{AB}$ and $\gamma_{a_1 \ldots a_{m+1}}{}^{AB}$ are always symmetric in their spinor indices.

Before we delve into the topic of pure spinors, we state without proof the following technical lemma that will be used later.

Lemma 2.2 *We have*

$$\begin{aligned} \gamma_{aA}{}^B \gamma_{b_1 \ldots b_p BD} \gamma_{cC}{}^D = (-1)^m \Big(& \gamma_{a b_1 \ldots b_p c AC} - (p+1) \gamma_{[a\, b_1 \ldots b_{p-1} AC}\, g_{\,b_p]c} \\ & - p g_{a[b_1}\, \gamma_{b_2 \ldots\, b_p]cAC} + p(p+1) g_{a[b_1}\, \gamma_{b_2 \ldots b_{p-1} AC}\, g_{\,b_p]c} \Big) \end{aligned}$$

In particular,

$$\gamma^a{}_A{}^B \gamma_{b_1 \ldots b_p BD} \gamma_{aC}{}^D = (-1)^{m+p} (2p - 2m - 1) \gamma_{b_1 \ldots b_p AC} \, .$$

2.2 Pure spinors

Let ξ^A be a non-zero spinor in $\mathfrak{S}$, and consider the map

$$\xi_a{}^A := \xi^B \gamma_{aB}{}^A : \mathfrak{V} \to \mathfrak{S} \, .$$

By the Clifford property (2.2), the kernel of $\xi_a{}^A : \mathfrak{V} \to \mathfrak{S}$ must be *totally null*, i.e. for any $X^a, Y^a \in \ker \xi_a{}^A : \mathfrak{V} \to \mathfrak{S}$, $g_{ab} X^a Y^b = 0$.

Definition 2.3 A non-zero spinor ξ^A is said to be *pure* if the kernel of $\xi_a{}^A : \mathfrak{V} \to \mathfrak{S}$ is m-dimensional.

Clearly, the purity property is invariant under rescaling of ξ^A.

Definition 2.4 A *null structure* or *γ-plane* on $\mathfrak{V}$ is a totally null m-dimensional vector subspace of $\mathfrak{V}$.

Remark 2.5 By identifying the spinor module $\mathfrak{S}$ with the $\mathcal{C}\ell(\mathfrak{V}, g)$-module $\wedge^\bullet \mathfrak{N}$ where $\mathfrak{N} \subset \mathfrak{V}$ is a null structure, it is clear that pure spinors exist and in fact span $\mathfrak{S}$. There is thus a one-to-one correspondence between γ-planes in $\mathfrak{V}$ and projective pure spinors.

Henceforth, we shall focus on the geometry arising from a fixed pure spinor ξ^A. By definition, ξ^A induces a filtration

$$\{0\} =: \mathfrak{V}^2 \subset \mathfrak{V}^1 \subset \mathfrak{V}^0 \subset \mathfrak{V}^{-1} \, . \tag{2.4}$$

where

$$\begin{aligned} &\mathfrak{S}^{\frac{m}{2}} := \langle \xi^A \rangle \, , && \mathfrak{S}^{\frac{m-2}{2}} := \operatorname{im} \xi_a{}^A : \mathfrak{V} \to \mathfrak{S} \, , && \\ &\mathfrak{V}^{-1} := \mathfrak{V} \, , && \mathfrak{V}^1 := \ker \xi_a{}^A : \mathfrak{V} \to \mathfrak{S} \, , && \mathfrak{V}^0 := (\mathfrak{V}^1)^\perp \, , \end{aligned}$$

where $\perp$ denotes the orthogonal complement with respect to g_{ab}, and one has an isomorphism[2]

$$\left(\mathfrak{V}^{-1} / \mathfrak{V}^1 \right) \otimes \mathfrak{S}^{\frac{m}{2}} \cong \mathfrak{S}^{\frac{m-2}{2}} \, , \tag{2.5}$$

of vector spaces. In particular, $\mathfrak{S}^{\frac{m-2}{2}}$ is an $(m+1)$-dimensional subspace of $\mathfrak{S}$, which must contain $\mathfrak{S}^{\frac{m}{2}}$, i.e.

$$\mathfrak{S}^{\frac{m}{2}} \subset \mathfrak{S}^{\frac{m-2}{2}} \, .$$

By invariance, this is identified with the 1-dimensional subspace $\mathfrak{V}^0 / \mathfrak{V}^1$, so that

$$(\mathfrak{V}^0 / \mathfrak{V}^1) \otimes \mathfrak{S}^{\frac{m}{2}} \cong \mathfrak{S}^{\frac{m}{2}} \, , \tag{2.6}$$

$$(\mathfrak{V}^{-1} / \mathfrak{V}^0) \otimes \mathfrak{S}^{\frac{m}{2}} \cong \mathfrak{S}^{\frac{m-2}{2}} / \mathfrak{S}^{\frac{m}{2}} \, , \tag{2.7}$$

Turning now to the dual picture, we set

$$\mathfrak{S}^{-\frac{m}{2}} := \mathfrak{S}^* \, , \qquad \mathfrak{S}^{-\frac{m-2}{2}} := \ker \xi^A : \mathbb{C} \leftarrow \mathfrak{S}^* \, , \qquad \mathfrak{S}^{-\frac{m-4}{2}} := \ker \xi_a{}^A : \mathfrak{V}^* \leftarrow \mathfrak{S}^* \, ,$$

so that, using the Clifford identity (2.2), we obtain the inclusion

$$\mathfrak{S}^{-\frac{m-4}{2}} \subset \mathfrak{S}^{-\frac{m-2}{2}} \subset \mathfrak{S}^{-\frac{m}{2}} \, .$$

---

[2] The use of factor $\mathfrak{S}^{\frac{m}{2}}$ on the LHS of (2.5) balances the 'degrees' on each side of (2.5), i.e. $-1 + \frac{m}{2} = \frac{m-2}{2}$.

Dual to (2.5), (2.6) and (2.7), we have

$$\mathfrak{V}^0 \cong \mathfrak{S}^{\frac{m}{2}} \otimes \left(\mathfrak{S}^{-\frac{m}{2}} / \mathfrak{S}^{-\frac{m-4}{2}} \right) , \tag{2.8}$$

$$\mathfrak{V}^0 / \mathfrak{V}^1 \cong \mathfrak{S}^{\frac{m}{2}} \otimes \left(\mathfrak{S}^{-\frac{m}{2}} / \mathfrak{S}^{-\frac{m-2}{2}} \right) , \tag{2.9}$$

$$\mathfrak{V}^1 \cong \mathfrak{S}^{\frac{m}{2}} \otimes \left(\mathfrak{S}^{-\frac{m-2}{2}} / \mathfrak{S}^{-\frac{m-4}{2}} \right) , \tag{2.10}$$

which can be expressed concretely as follows.

Lemma 2.6 *Let V^a be a non-zero vector in $\mathfrak{V}$. Then*

- *V^a is an element of $\mathfrak{V}^0$ if and only if $V^a = \xi^{aA} v_A$ for some non-zero $v_A \in \mathfrak{S}^{-\frac{m}{2}} / \mathfrak{S}^{-\frac{m-4}{2}}$;*
- *V^a is an element of $\mathfrak{V}^1$ if and only if $V^a = \xi^{aA} v_A$ for some non-zero $v_A \in \mathfrak{S}^{-\frac{m-2}{2}} / \mathfrak{S}^{-\frac{m-4}{2}}$.*

Proposition 2.7 *A spinor ξ^A is pure if and only if it satisfies*

$$\xi^{aC} \xi_a{}^D = - \xi^C \xi^D . \tag{2.11}$$

The factor of -1 can be determined using Lemma 2.2 and Proposition 2.8, which is the more familiar algebraic characterisation of pure spinors due to Cartan [Car81].

Proposition 2.8 (Cartan [Car81]**)** *A non-zero spinor $\xi^{A'}$ is pure if and only if it satisfies*

$$\begin{aligned} & \gamma_{a_1 \dots a_p AB} \xi^A \xi^B = 0 , \quad \text{for all } p < m,\ m - p \equiv 0, 3 \pmod 4, \\ & \gamma_{a_1 \dots a_m AB} \xi^A \xi^B \neq 0 . \end{aligned} \tag{2.12}$$

We shall refer to both equations (2.11) and (2.12) as the *purity conditions* of a spinor ξ^A. These are vacuous when $m \leq 2$, i.e. all spinors are pure when $m \leq 2$.

The only non-vanishing irreducible components of the tensor product $\xi^A \xi^B$ is thus the m-form

$$\phi_{a_1 \dots a_m} := \gamma_{a_1 \dots a_m AB} \xi^A \xi^B ,$$

which can be seen to annihilate $\mathfrak{V}^0$. It is *simple* (or *decomposable*) in the sense that

$$\phi_{a_1 \dots a_m} = \xi_{a_1}{}^{A_1} \dots \xi_{a_m}{}^{A_m} \varepsilon_{A_1 \dots A_m} \quad \in \quad \wedge^m \mathfrak{V}^1 ,$$

for some $\varepsilon_{A_1 \dots A_m} \in \wedge^m \left(\mathfrak{S}^{-\frac{m-2}{2}} / \mathfrak{S}^{-\frac{m-4}{2}} \right)$. Its Hodge dual annihilates $\mathfrak{V}^1$ and takes the form

$$(*\phi)_{a_1 \dots a_{m+1}} = \xi_{a_1}{}^{A_1} \dots \xi_{a_{m+1}}{}^{A_{m+1}} \varepsilon_{A_1 \dots A_{m+1}} \quad \in \quad \wedge^{m+1} \mathfrak{V}^0 ,$$

for some $\varepsilon_{A_1 \dots A_{m+1}} \in \wedge^{m+1} \left(\mathfrak{S}^{-\frac{m}{2}} / \mathfrak{S}^{-\frac{m-4}{2}} \right)$.

Splitting For clarity, we can choose a spinor η_A dual to ξ^A, i.e. η_A belongs to $\mathfrak{S}^{-\frac{m}{2}}$, but not to $\mathfrak{S}^{-\frac{m-2}{2}}$, and satisfying $\xi^A \eta_A = -\frac{1}{2}$. This distinguishes a one-dimensional subspace $\langle \eta_A \rangle =: \mathfrak{S}_{-\frac{m}{2}} \subset \mathfrak{S}^{-\frac{m}{2}}$ complementary to $\mathfrak{S}^{-\frac{m-2}{2}}$. One associates to η_A the map $\eta_{aA} := \eta_B \gamma_{aA}{}^B : \mathfrak{V} \to \mathfrak{S}^*$. Now, it is straightforward to see that the map

$$I_A^B := \eta_{aA} \xi^{aB} + \eta_A \xi^B : \mathfrak{S} \to \mathfrak{S} \tag{2.13}$$

is idempotent, has trace $I_A^A = m$, and is degenerate on $\mathfrak{S}^{\frac{m}{2}}$, i.e. $\xi^A I_A^B = 0$. Thus, I_A^B is the identity element on $\mathfrak{S}_{\frac{m-2}{2}}$, the complement of $\mathfrak{S}^{\frac{m}{2}}$ in $\mathfrak{S}^{\frac{m-2}{2}}$. In fact, η_A is pure with associated γ-plane $\mathfrak{V}_{-1}$ dual to $\mathfrak{V}_1 := \mathfrak{V}^1$. We can then write

$$\mathfrak{V} = \mathfrak{V}_{-1} \oplus \mathfrak{V}_0 \oplus \mathfrak{V}_1 \,, \tag{2.14}$$

where

$$\mathfrak{S}_{-\frac{m-2}{2}} := \{\operatorname{im} \eta_{aA} : \mathfrak{V} \to \mathfrak{S}^*\} \cap \{\ker \xi^A : \mathbb{C} \leftarrow \mathfrak{S}^*\} \,, \qquad \mathfrak{V}_{-1} := \ker \eta^a{}_A : \mathfrak{V} \to \mathfrak{S} \,.$$

For future use, we set

$$u^a := 2\, \xi^{aA} \eta_A \,, \tag{2.15}$$

which can be seen to square to -1 and satisfies $u^a \xi_a{}^A = \xi^A$ and $u^a \eta_{aA} = \eta_A$. The metric tensor g_{ab} together with a choice of u^a determines a symmetric bilinear form

$$h_{ab} = g_{ab} + u_a u_b \,, \tag{2.16}$$

on $\mathfrak{V}_0$, i.e. $h_{ab} u^a = 0$, and $h_a{}^c h_c{}^b = h_a{}^b$, and $h_a{}^a = n - 1$.

Intersection of γ-planes

Proposition 2.9 *Any spinor β^A in $\mathfrak{S}^{\frac{m-2}{2}}$ is a pure spinor. Further, if β^A is not in $\mathfrak{S}^{\frac{m}{2}}$, the respective γ-planes defined by β^A and ξ^A intersect in a totally null $(m-1)$-plane. This is algebraically equivalent to β^A and ξ^A satisfying*

$$\beta^{aA} \xi_a{}^B = \beta^A \xi^B - 2 \xi^A \beta^B = -\beta^{(A} \xi^{B)} + 3 \beta^{[A} \xi^{B]} \,,$$

or equivalently,

$$\begin{aligned} \gamma_{a_1 \ldots a_p AB} \beta^A \xi^B &= 0 \,, & &\text{for all } p < m-2, \\ \gamma_{a_1 \ldots a_{m-2} AB} \beta^A \xi^B &\neq 0 \,. \end{aligned}$$

Further, the γ-planes of two distinct projective spinors β^A and ρ^A in $\mathfrak{S}^{\frac{m-2}{2}}$ not in $\mathfrak{S}^{\frac{m}{2}}$ intersect in a totally null $(m-2)$-plane. This is algebraically equivalent to β^A and ρ^A satisfying

$$\beta^{a(A} \rho_a{}^{B)} = -\beta^{(A} \rho^{B)}$$

or equivalently,

$$\begin{aligned} \gamma_{a_1 \ldots a_p AB} \beta^A \rho^B &= 0 \,, & &\text{for all } p < m-3, \\ \gamma_{a_1 \ldots a_{m-3} AB} \beta^A \rho^B &\neq 0 \,. \end{aligned}$$

Remark 2.10 We omit the proof of the above proposition, which was essentially given by Cartan [Car81]. The second part of the proposition is a restatement of a standard result that the sum of two pure spinors is again pure if and only if their corresponding γ-planes intersect in an $(m-2)$-planes.

2.3 Spinorial description of the Lie algebra $\mathfrak{so}(2m+1,\mathbb{C})$

Filtration We can extend the above calculus in the classification of the Lie algebra $\mathfrak{g} = \mathfrak{so}(2m+1,\mathbb{C})$, or equivalently to the space of 2-forms $\wedge^2\mathfrak{V}$. For $\phi_{ab} \in \mathfrak{g}$, define

$$ {}^{\mathfrak{g}}\Pi_{-2}(\phi) := \xi^{a[A}\xi^{bB}\phi_{ab}\xi^{C]} \,, \quad {}^{\mathfrak{g}}\Pi_{-1}(\phi) := \xi^{aA}\xi^{bB}\phi_{ab} \,, \quad {}^{\mathfrak{g}}\Pi_{0}(\phi) := \xi^{a[A}\phi_{ab}\xi^{B]} \,, \quad {}^{\mathfrak{g}}\Pi_{1}(\phi) := \xi^{aA}\phi_{ab} \,, $$

and set

$$ \mathfrak{g}^i := \{\phi_{ab} \in \mathfrak{g} : {}^{\mathfrak{g}}\Pi_{i-1}(\phi) = 0\} \,. $$

turns $\mathfrak{g}$ into a filtered Lie algebra, i.e. there is a filtration

$$ \{0\} =: \mathfrak{g}^3 \subset \mathfrak{g}^2 \subset \mathfrak{g}^1 \subset \mathfrak{g}^0 \subset \mathfrak{g}^{-1} \subset \mathfrak{g}^{-2} := \mathfrak{g} \,. $$

on $\mathfrak{g}$, with which the Lie bracket on $\mathfrak{g}$ is compatible.

Proposition 2.11 *The Lie subalgebra $\mathfrak{g}^0$ is the stabilizer of ξ^A, i.e.*

$$ \phi_{ab}\xi^{abA} \propto \xi^A \,. $$

Proof. We only need to show $\phi_{ab}\xi^{aA}\xi^{bB} = 0$ if and only if $\phi_{ab}\xi^{ab[A}\xi^{B]} = 0$. To see this, we use the identities

$$ \phi_{ab}\xi^{aA}\xi^{bB}\gamma^{c}{}_{A}{}^{C}\gamma_{cB}{}^{D} = -\phi_{ab}\left(\xi^{aC}\xi^{bD} + 4\xi^{ab[C}\xi^{D]}\right) \,, \tag{2.17} $$

$$ \phi_{ab}\xi^{abA}\gamma^{c}{}_{A}{}^{C}\xi_{c}{}^{D} = -\phi_{ab}\left(\xi^{abC}\xi^{D} + 4\xi^{aC}\xi^{bD}\right) \,. \tag{2.18} $$

The first of these gives ⇒ immediately, while the second gives ⇐ by rewriting the LHS as $\phi_{ab}\xi^{abA} = \phi\xi^A$ for some ϕ. ☐

Associated graded vector space We shall refine the filtration $\{\mathfrak{g}^i\}$ on $\mathfrak{g}$ in the moment. For this purpose, we define

$$ \xi_{ab}{}^{A} := \xi^{B}\gamma_{abB}{}^{A} : \wedge^2\mathfrak{V} \to \mathfrak{S} \,, \qquad \mathfrak{S}^{\frac{m-4}{2}} := \operatorname{im}\xi_{ab}{}^{A} : \wedge^2\mathfrak{V} \to \mathfrak{S} \,. $$

We note that one must have

$$ \mathfrak{S}^{\frac{m}{2}} \subset \mathfrak{S}^{\frac{m-2}{2}} \subset \mathfrak{S}^{\frac{m-4}{2}} \,. $$

The first inclusion has already been shown, while the second one can be proved along the same line: we have $0 = \xi_{ab}{}^{A}\eta_A = \xi_a{}^{B}\gamma_{bB}{}^{A}\eta_A + g_{ab}\xi^A\eta_A$. Now, contracting the above equation with any $V^a \in \mathfrak{V}^1$ and $W^b \in \mathfrak{V}^{-1}$ such that $V^aW_a \neq 0$, yields $0 = V^aW_a\xi^{A'}\eta_{A'}$, and contracting the above equation with any non-null $V^a \in \mathfrak{V}^0$ gives the inclusion. It is not too difficult to show

$$ \left(\mathfrak{g}^{-2}/\mathfrak{g}^{-1}\right) \otimes \mathfrak{S}^{\frac{m}{2}} \cong \mathfrak{S}^{\frac{m-4}{2}}/\mathfrak{S}^{\frac{m-2}{2}} \,, \qquad \left(\mathfrak{g}^{-1}/\mathfrak{g}^{0}\right) \otimes \mathfrak{S}^{\frac{m}{2}} \cong \mathfrak{S}^{\frac{m-2}{2}}/\mathfrak{S}^{\frac{m}{2}} \,. $$

and dually,

$$ \mathfrak{g}^{2} \cong \mathfrak{S}^{\frac{m}{2}} \otimes \left(\mathfrak{S}^{-\frac{m-4}{2}}/\mathfrak{S}^{-\frac{m-6}{2}}\right) \,, \qquad \mathfrak{g}^{-1} \cong \mathfrak{S}^{\frac{m}{2}} \otimes \left(\mathfrak{S}^{-\frac{m-2}{2}}/\mathfrak{S}^{-\frac{m-4}{2}}\right) \,, $$

where $\mathfrak{S}^{\frac{m-6}{2}} := \ker\xi_{ab}{}^{A} : \wedge^2\mathfrak{V} \leftarrow \mathfrak{S}^*$.

The map ${}^{\mathfrak{g}}\Pi_0(\phi)$ is not saturated with symmetries, and can be split further in terms of

$$ {}^{\mathfrak{g}}\Pi_0^0(\phi) := \xi^{abA}\phi_{ab} \,, \qquad {}^{\mathfrak{g}}\Pi_0^1(\phi) := \xi^{a[A}\phi_{ab}\xi^{B]} + \frac{1}{n-1}\xi^{acC}\phi_{ac}\gamma_{bC}{}^{[A}\xi^{B]} \,. $$

It follows that $\mathfrak{g}^0/\mathfrak{g}^1 \cong \mathfrak{g}_0^0 \oplus \mathfrak{g}_0^1$ where we have defined the vector subspaces

$$ \mathfrak{g}_0^0 := \left\{\phi_{ab} \in \mathfrak{g}^0 : {}^{\mathfrak{g}}\Pi_0^1(\phi) = 0\right\}/\mathfrak{g}^1 \,, \qquad \mathfrak{g}_0^1 := \left\{\phi_{ab} \in \mathfrak{g}^0 : {}^{\mathfrak{g}}\Pi_0^0(\phi) = 0\right\}/\mathfrak{g}^1 \,. \tag{2.19} $$

For later use, we set $\mathfrak{g}_i^0 := \mathfrak{g}^i/\mathfrak{g}^{i+1}$ for $i = \pm 1, \pm 2$.

Splitting The splitting (2.14) extends to tensor products of $\mathfrak{V}$ and $\mathfrak{V}^*$, and in particular, the Lie algebra $\mathfrak{g}$ admits the structure of $|2|$-graded Lie algebra

$$\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2 \,, \qquad\qquad [\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j} \,,$$

for all i, j, where $\mathfrak{g}_i \subset \mathfrak{g}^i$ is complementary to $\mathfrak{g}^{i-1}$ in $\mathfrak{g}^i$, and we can make the identification

$$\mathfrak{g}_{-2} \cong \wedge^2 \mathfrak{V}_{-1} \,, \qquad \mathfrak{g}_{-1} \cong \mathfrak{V}_0 \otimes \mathfrak{V}_{-1} \,, \qquad \mathfrak{g}_0 \cong \mathfrak{V}_{-1} \otimes \mathfrak{V}_1 \,, \qquad \mathfrak{g}_1 \cong \mathfrak{V}_0 \otimes \mathfrak{V}_1 \,, \qquad \mathfrak{g}_2 \cong \wedge^2 \mathfrak{V}_1 \,.$$

Using ξ^A, η_A and u^a defined by (2.15), one can give a spinorial decomposition of any element ϕ_{ab} of $\mathfrak{g}$ as

$$\begin{aligned} \phi_{ab} &= \xi_a{}^A \xi_b{}^B \phi_{AB} - 2\xi_{[a}{}^A u_{b]} \phi_A + 2\xi_{[a}{}^A \eta_{b]B} \phi_A{}^B - 2\eta_{[a\,A} u_{b]} \phi^A + \eta_{aA} \eta_{bB} \phi^{AB} \\ &\in \quad \mathfrak{g}_2 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_{-2} \,, \end{aligned}$$

where $\phi_{AB} = \phi_{[AB]}$, $\phi^{AB} = \phi^{[AB]}$, $\phi_A{}^B$, ϕ_A and ϕ^A are all elements of $\mathfrak{S}_{\frac{m-2}{2}}$ and $\mathfrak{S}_{-\frac{m-2}{2}}$, and tensor products thereof.

The Lie subalgebra $\mathfrak{g}_0$ is isomorphic to $\mathfrak{gl}(m, \mathbb{C})$, and thus splits futher as $\mathfrak{g}_0 = \mathfrak{z}_0 \oplus \mathfrak{sl}_0$ where $\mathfrak{z}_0$ is the center $\mathfrak{g}_0$ and $\mathfrak{sl}_0 = \mathfrak{sl}(m, \mathbb{C})$ is the semisimple part of $\mathfrak{g}_0$. The center is spanned by the *grading element*

$$E_{ab} := -2\xi_{[a}{}^A \eta_{b]A} = -2\xi_a{}^A \eta_{bA} + g_{ab} \,,$$

and from the identity $\xi^{aA} E_a{}^b = 2\xi^A \xi^{bB} \eta_B + \xi^{bB}$, one can easily deduce that $E_a{}^b$ has eigenvalues i on $\mathfrak{V}_i$. Its image in the Clifford algebra is given by $E_A{}^B := -\frac{1}{4} E_{ab} \gamma^{ab}{}_A{}^B$, and a straightforward computation yields $E_B{}^A \xi^B = \frac{m}{2} \xi^A$, and similarly for the dual action.

For consistency with reference [TC12b], we set

$$\omega_{ab} := E_{ab} \,. \tag{2.20}$$

Then an element of $\mathfrak{g}_0$ can be expressed as

$$\phi_{ab} = \phi \, \omega_{ab} + 2 \, \xi_{[a}{}^A \eta_{b]B} \phi_A{}^B$$

for some complex ϕ and tracefree $\phi_A{}^B$, i.e. $\phi_A{}^B I_B^A = 0$ where I_A^B is the identity element defined by (2.13).

The above decomposition thus gives us an explicit way of describind the vector spaces (in fact $\mathfrak{p}$-modules) $\mathfrak{g}_i^j$ defined in (2.19) and after. To be precise, we have isomorphisms

$$\mathfrak{g}_{\pm 2}^0 \cong \breve{\mathfrak{g}}_{\pm 2}^0 := \mathfrak{g}_{\pm 2} \,, \qquad \mathfrak{g}_{\pm 1}^0 \cong \breve{\mathfrak{g}}_{\pm 1}^0 := \mathfrak{g}_{\pm 1} \,, \qquad \mathfrak{g}_0^0 \cong \breve{\mathfrak{g}}_0^0 := \mathfrak{z}_0 \,, \qquad \mathfrak{g}_0^1 \cong \breve{\mathfrak{g}}_0^1 := \mathfrak{sl}_0 \,.$$

between $\mathfrak{p}$-modules and $\mathfrak{g}_0$-modules.

The additional information needed to describe $\mathfrak{g}$ in terms of irreducible $\mathfrak{p}$-modules comes from the (non-trivial) action of the nilpotent part of $\mathfrak{p}$, or more simply $\mathfrak{g}_1$ on the $\mathfrak{g}_0$-modules $\mathfrak{g}_i^j$. This orbit of $\mathfrak{p}$ on $\mathfrak{g}$ can be encoded by means of the directed graph (a 'Penrose diagram')

$$\begin{tikzcd} & & \mathfrak{g}_0^0 \arrow[dr] & & \\ \mathfrak{g}_2^0 \arrow[r] & \mathfrak{g}_1^0 \arrow[ur] \arrow[dr] & & \mathfrak{g}_{-1}^0 \arrow[r] & \mathfrak{g}_{-2}^0 \\ & & \mathfrak{g}_0^1 \arrow[ur] & & \end{tikzcd}$$

where an arrow from $\mathfrak{g}_i^j$ to $\mathfrak{g}_{i-1}^k$ simply means that $\mathfrak{g}_1 \cdot \breve{\mathfrak{g}}_i^j \subset \breve{\mathfrak{g}}_{i-1}^k$, or dually that

$$\ker {}^{\mathfrak{g}}\Pi_0^j \subset \ker {}^{\mathfrak{g}}\Pi_{-1}^0 \,, \qquad\qquad j = 0, \, 1 \,. \tag{2.21}$$

This is of course a rather trivial example, but such Penrose diagrams will be generalised to more complicated $\mathfrak{g}$-modules in sections 3 and 4.

Parabolic Lie subgroups We can translate the description of the Lie algebra $\mathfrak{g}$ and its parabolic Lie subalgebra $\mathfrak{p}$ to the Lie group level. We shall take G to be the complex special orthogonal group $\mathrm{SO}(2m+1,\mathbb{C})$. Then the *parabolic Lie subgroup* P of G stabilising the pure spinor ξ^A can simply be obtained by exponientiating $\mathfrak{p}$. In particular, it admits a Levy decomposition $P = G_0 \times P_+$ where G_0 is isomorphic to the complex general linear group $\mathrm{GL}(m,\mathbb{C})$ and P_+ is the abelian Lie group generated by $\mathfrak{g}_1 \oplus \mathfrak{g}_2$. The spinor calculus developed here is then manifestly P-invariant.

Denote the null (or isotropic) Grassmanian $\mathrm{Gr}_m(\mathfrak{V}, g)$ of γ-planes in $\mathfrak{V}$. By definition, $\mathrm{Gr}_m(\mathfrak{V}, g)$ can also be described as the space of projective pure spinors of $(\mathfrak{V}, g)$, i.e. a compact complex subvariety of $\mathbb{P}\mathfrak{S}$ defined by the purity conditions (2.12). Since the action of $\mathrm{SO}(2m+1,\mathbb{C})$ is transitive on $\mathrm{Gr}_m(\mathfrak{V}, g)$ and P stabilises a point in $\mathrm{Gr}_m(\mathfrak{V}, g)$, the space of projective pure spinors of $(\mathfrak{V}, g)$ is simply the $\frac{1}{2}m(m+1)$-dimensional homogeneous space G/P. When $m = 1, 2$, this space is isomorphic to the complex projective space $\mathbb{CP}^{\frac{1}{2}m(m+1)}$. See [FH91, HS95] for more details.

Remark 2.12 By Proposition (2.9), each (non-projective) pure spinor defines an $(m+1)$-dimensional vector subspace of non-projective spinors. Therefore, to each point in $\mathrm{Gr}_m(\mathfrak{V}, g)$ corresponds an m-dimensional complex projective space $\mathbb{CP}^m$ in $\mathrm{Gr}_m(\mathfrak{V}, g)$. This is trivial when $m = 1$, and when $m = 2$, this is simply the hyperplane dual to a point in $\mathbb{CP}^3$.

2.4 Generalisation to $\mathfrak{so}(2m+1,\mathbb{C})$-modules

The filtration (2.4) of $\mathfrak{p}$-modules defined by the projective pure spinor $[\xi^A]$ induces a filtration

$$\{0\} = \mathfrak{R}^{p+1} \subset \mathfrak{R}^{p} \subset \mathfrak{R}^{p-1} \subset \ldots \subset \mathfrak{R}^{-p+1} \subset \mathfrak{R}^{-p} = \mathfrak{R}$$

of $\mathfrak{p}$-modules on $\mathfrak{R} := \bigotimes^p \mathfrak{V}$ for any $p \geq 0$, where

$$\mathfrak{R}^j := \sum_{i_1+\ldots+i_p=j} \mathfrak{V}^{i_1} \otimes \ldots \otimes \mathfrak{V}^{i_p} \,.$$

We can then say that $\xi^{A'}$ is a *k-fold principal spinor* of a tensor $T \in \mathfrak{R}$, if T lies in $\mathfrak{R}^{-p+k}$, but not in $\mathfrak{R}^{-p+k+1}$. However, as in the even-dimensional, it is best to replace the concept of principal spinors with that of P-invariant filtrations on $\mathfrak{g}$-modules.

The filtration on the spinor module $\mathfrak{S}$ can be described as follows. Given a (projective) pure spinor ξ^A, we can define the maps

$$\xi_{a_1\ldots a_k}{}^A := \xi^B \gamma_{a_1\ldots a_{2k+1}B}{}^A : \wedge^k\mathfrak{V} \to \mathfrak{S} \,.$$

By Hodge duality, it is enough to consider values of k ranging from 0 to m, with the understanding that $\xi^A : \wedge^0\mathfrak{V} \cong \mathbb{C} \to \mathfrak{S}$. An argument identical to the discussion so far then tells us that ξ^A induces filtration

$$\mathfrak{S}^{\frac{m}{2}} \subset \mathfrak{S}^{\frac{m-2}{2}} \subset \ldots \subset \mathfrak{S}^{-\frac{m-2}{2}} \subset \mathfrak{S}^{-\frac{m}{2}} = \mathfrak{S} \,,$$

on $\mathfrak{S} \cong \mathfrak{S}^*$, where we have defined

$$\mathfrak{S}^{\frac{m-2k}{2}} := \mathrm{im}\,\xi_{a_1\ldots a_k}{}^A : \wedge^k\mathfrak{V} \to \mathfrak{S} \,, \qquad \mathfrak{S}^{-\frac{m-2k}{2}} = \ker\xi_{a_1\ldots a_k}{}^A : \wedge^k\mathfrak{V}^* \leftarrow \mathfrak{S}^* \,.$$

Further, we can choose subspaces $\mathfrak{S}_i \subset \mathfrak{S}^i$ such that $\mathfrak{S}^i = \mathfrak{S}_i \oplus \mathfrak{S}^{i+1}$ such that

$$\mathfrak{S} = \mathfrak{S}_{\frac{m}{2}} \oplus \mathfrak{S}_{\frac{m-2}{2}} \oplus \ldots \oplus \mathfrak{S}_{-\frac{m-2}{2}} \oplus \mathfrak{S}_{-\frac{m}{2}} \,.$$

The grading element E_{ab} in $\mathfrak{z}_0$ and the spanning element u_a of $\mathfrak{V}_0$ have eigenvalues $\frac{m-2k}{2}$ and $(-1)^k$ respecively on $\mathfrak{S}_{\frac{m-2k}{2}}$.

Remark 2.13 In the form description of the spinor representation $\mathfrak{S}$, we can make the identification $\mathfrak{S}_{\frac{m-2i}{2}} \cong \wedge^{m-i}\mathfrak{N}$.

The generalisation of isomorphisms (2.10) and are given by

$$\begin{aligned} \wedge^k \mathfrak{V}^1 &\cong \mathfrak{S}^{\frac{m}{2}} \otimes \left(\mathfrak{S}^{-\frac{m-2k}{2}} / \mathfrak{S}^{-\frac{m-2k-2}{2}} \right) , \qquad k = 0 , \ldots , m-1 , \\ \wedge^m \mathfrak{V}^1 &\cong \wedge^m \mathfrak{V}^1 \otimes \mathfrak{V}^0 \cong \mathfrak{S}^{\frac{m}{2}} \otimes \mathfrak{S}^{\frac{m}{2}} . \end{aligned}$$

2.5 Real pure spinors

As in even dimensions, the above calculus applies equally in the case where $(\mathfrak{V}, g)$ is a real vector space equipped with a non-degenerate bilinear form of signature $(m, m+1)$ or $(m+1, m)$, and our choice of pure spinor is *real*, so that its associated γ-plane is a real totally null subspace of $\mathfrak{V}$. The description of its stabiliser $\mathfrak{p}$ as a real parabolic Lie subalgebra of $\mathfrak{so}(m, m)$, and similarly for its Lie group P as a subgroup of the connected identity component of $\mathrm{SO}(m, m)$.

If the bilinear form g_{ab} is now taken to be positive definite, one obtains an $\mathrm{SO}(2m+1)$-compatible CR structure, which is none other than an odd-dimensional analogue of a Hermitian structure.

If the bilinear form g_{ab} is of Lorentzian signature, i.e. $(1, 2m)$ or $(2m, 1)$, the resulting geometry is that of a *Robinson structure* [NT02, Tra02] – this is treated in more details in [TC].

3 Algebraic classification of curvature tensors

We now apply the ideas developed in the previous section to the classifications of the spaces $\mathfrak{F}$, $\mathfrak{A}$ and $\mathfrak{C}$ of tracefree symmetric 2-tensors, of tensors with Cotton-York symmetries and of tensors with Weyl symmetries respectively, i.e.

$$\begin{aligned} \mathfrak{A} &:= \{ A_{abc} \in \otimes^3 \mathfrak{V} : A_{abc} = A_{a[bc]} \, , A_{[abc]} = 0 \, , A^a{}_{ac} = 0 \} \, , \\ \mathfrak{F} &:= \{ \Phi_{ab} \in \otimes^2 \mathfrak{V}^* : \Phi_{ab} = \Phi_{(ab)} \, , \Phi_c{}^c = 0 \} \, , \\ \mathfrak{C} &:= \{ C_{abcd} \in \otimes^4 \mathfrak{V} : C_{abcd} = C_{[ab][cd]} \, , C_{[abc]d} = 0 \, , C^a{}_{bad} = 0 \} \, . \end{aligned}$$

From a representation theoretic viewpoint, the $\mathfrak{p}$-invariant decomposition of tensors follows the same algorithm as in the even-dimensional case. This is explained in details in [TC12b] – see also [ČS09b]. One can also describe these irreducibles explictly in terms of kernels of certain maps projecting into them,[3] but compared to the even-dimensional case, these maps are somewhat a little more 'baroque'.

3.1 Classification of the tracefree Ricci tensor

Proposition 3.1 *The space $\mathfrak{F}$ of tracefree symmetric 2-tensors admits a filtration*

$$\{0\} =: \mathfrak{F}^3 \subset \mathfrak{F}^2 \subset \mathfrak{F}^1 \subset \mathfrak{F}^0 \subset \mathfrak{F}^{-1} \subset \mathfrak{F}^{-2} := \mathfrak{F} \, ,$$

of $\mathfrak{p}$-modules.

The associated graded $\mathfrak{p}$-module $\mathrm{gr}(\mathfrak{F}) = \bigoplus_{i=-2}^{2} \mathrm{gr}_i(\mathfrak{F})$, where $\mathrm{gr}_i(\mathfrak{F}) := \mathfrak{F}^i / \mathfrak{F}^{i+1}$ splits into a direct sum of irreducible $\mathfrak{p}$-modules

$$\mathrm{gr}_{\pm 2}(\mathfrak{F}) = \mathfrak{F}^0_{\pm 2} \, , \qquad \mathrm{gr}_{\pm 1}(\mathfrak{F}) = \mathfrak{F}^0_{\pm 1} \, , \qquad \mathrm{gr}_0(\mathfrak{F}) = \mathfrak{F}^0_0 \oplus \mathfrak{F}^1_0 \, ,$$

where the $\mathfrak{F}^j_i$ are given in Table 3 of appendix D. Further,

$$\mathfrak{F}^j_0 := \{ \Phi_{ab} \in \mathfrak{F}^0 : {}^{\mathfrak{F}}\Pi^j_0(\Phi) = 0 \} / \mathfrak{F}^1 \, ,$$

---

[3] These are denoted ${}^{\mathfrak{F}}\Pi^j_i$, ${}^{\mathfrak{A}}\Pi^j_i$ and ${}^{\mathfrak{C}}\Pi^j_i$ below.

where the maps ${}^{\mathfrak{F}}\Pi^j_0$ *are defined in A.*

Finally, the $\mathfrak{p}$*-orbit structure of this decomposition can be expressed by means of the graph*

$$\begin{tikzcd}
& & \mathfrak{F}^1_0 \arrow[dr] & & \\
\mathfrak{F}^0_2 \arrow[r] & \mathfrak{F}^0_1 \arrow[ur] \arrow[dr] & & \mathfrak{F}^0_{-1} \arrow[r] & \mathfrak{F}^0_{-2} \\
& & \mathfrak{F}^0_0 \arrow[ur] & &
\end{tikzcd}$$

where an arrow from $\mathfrak{F}^j_i$ *to* $\mathfrak{F}^k_{i-1}$ *for some* i, j, k *implies that* $\breve{\mathfrak{F}}^j_i \subset \mathfrak{g}_1 \cdot \breve{\mathfrak{F}}^k_{i-1}$ *for any choice of irreducible* $\mathfrak{g}_0$*-modules* $\breve{\mathfrak{F}}^j_i$ *and* $\breve{\mathfrak{F}}^k_{i-1}$ *isomorphic to* $\mathfrak{F}^j_i$ *and* $\mathfrak{F}^k_{i-1}$ *respectively.*

3.2 Classification of the Cotton-York tensor

Proposition 3.2 *The space* $\mathfrak{A}$ *of tensors with Cotton-York symmetries admits a filtration*

$$\{0\} =: \mathfrak{A}^4 \subset \mathfrak{A}^3 \subset \mathfrak{A}^2 \subset \mathfrak{A}^1 \subset \mathfrak{A}^0 \subset \mathfrak{A}^{-1} \subset \mathfrak{A}^{-2} \subset \mathfrak{A}^{-3} := \mathfrak{A}\,,$$

of $\mathfrak{p}$*-modules.*

The associated graded $\mathfrak{p}$*-module* $\mathrm{gr}(\mathfrak{A}) = \bigoplus_{i=-3}^{3} \mathrm{gr}_i(\mathfrak{A})$*, where* $\mathrm{gr}_i(\mathfrak{A}) := \mathfrak{A}^i/\mathfrak{A}^{i+1}$ *splits into a direct sum of irreducible* $\mathfrak{p}$*-modules*

$$\begin{aligned}
\mathrm{gr}_{\pm 3}(\mathfrak{A}) &= \mathfrak{A}^0_{\pm 3}\,, & \mathrm{gr}_{\pm 2}(\mathfrak{A}) &= \mathfrak{A}^0_{\pm 2} \oplus \mathfrak{A}^1_{\pm 2}\,, \\
\mathrm{gr}_{\pm 1}(\mathfrak{A}) &= \mathfrak{A}^0_{\pm 1} \oplus \mathfrak{A}^1_{\pm 1} \oplus \mathfrak{A}^2_{\pm 1} \oplus \mathfrak{A}^3_{\pm 1}\,, & \mathrm{gr}_0(\mathfrak{A}) &= \mathfrak{A}^0_0 \oplus \mathfrak{A}^1_0 \oplus \mathfrak{A}^2_0\,,
\end{aligned}$$

where the $\mathfrak{A}^j_i$ *are given in Table 4 of appendix D. Further,*

$$\mathfrak{A}^j_i = \{A_{abc} \in \mathfrak{A}^i : {}^{\mathfrak{A}}\Pi^k_i(A) = 0 \textit{ for all } k \neq j\}/\mathfrak{A}^{i+1}\,,$$

for $|i| \leq 2$*, where the maps* ${}^{\mathfrak{C}}\Pi^j_i$ *are defined in appendix A.*

Finally, the $\mathfrak{p}$*-orbit structure of this decomposition can be expressed by means of the graph*

$$\begin{tikzcd}
& \mathfrak{A}^1_2 \arrow[r] \arrow[ddr] & \mathfrak{A}^3_1 \arrow[r] & \mathfrak{A}^1_0 \oplus \mathfrak{A}^2_0 \arrow[r] \arrow[ddr] & \mathfrak{A}^3_{-1} \arrow[r] & \mathfrak{A}^1_{-2} \arrow[dr] & \\
\mathfrak{A}^0_3 \arrow[ur] \arrow[dr] & & & & & & \mathfrak{A}^0_{-3} \\
& \mathfrak{A}^0_2 \arrow[r] & \mathfrak{A}^0_1 \oplus \mathfrak{A}^1_1 \oplus \mathfrak{A}^2_1 \arrow[r] \arrow[uur] & \mathfrak{A}^0_0 \arrow[r] & \mathfrak{A}^0_{-1} \oplus \mathfrak{A}^1_{-1} \oplus \mathfrak{A}^2_{-1} \arrow[r] \arrow[uur] & \mathfrak{A}^0_{-2} \arrow[ur] &
\end{tikzcd}$$

with the proviso that when $m = 2$*,* $\mathfrak{A}^2_{\pm 1}$ *does not occur. Here an arrow from* $\mathfrak{A}^j_i$ *to* $\mathfrak{A}^k_{i-1}$ *for some* i, j, k *implies that* $\breve{\mathfrak{A}}^j_i \subset \mathfrak{g}_1 \cdot \breve{\mathfrak{A}}^k_{i-1}$ *for any choice of irreducible* $\mathfrak{g}_0$*-modules* $\breve{\mathfrak{A}}^j_i$ *and* $\breve{\mathfrak{A}}^k_{i-1}$ *isomorphic to* $\mathfrak{A}^j_i$ *and* $\mathfrak{A}^k_{i-1}$ *respectively.*

3.3 Classification of the Weyl tensor

Proposition 3.3 *The space* $\mathfrak{C}$ *of tensors with Weyl symmetries admits a filtration*

$$\{0\} =: \mathfrak{C}^5 \subset \mathfrak{C}^4 \subset \mathfrak{C}^3 \subset \mathfrak{C}^2 \subset \mathfrak{C}^1 \subset \mathfrak{C}^0 \subset \mathfrak{C}^{-1} \subset \mathfrak{C}^{-2} \subset \mathfrak{C}^{-3} \subset \mathfrak{C}^{-4} := \mathfrak{C}\,,$$

of $\mathfrak{p}$*-modules* $\mathfrak{C}^i$*.*

The associated graded $\mathfrak{p}$*-module* $\mathrm{gr}(\mathfrak{C}) = \bigoplus_{i=-4}^{4} \mathrm{gr}_i(\mathfrak{C})$*, where* $\mathrm{gr}_i(\mathfrak{C}) := \mathfrak{C}^i/\mathfrak{C}^{i+1}$ *splits into a direct sum of irreducible* $\mathfrak{p}$*-modules*

$$\begin{aligned} &\mathrm{gr}_{\pm 4}(\mathfrak{C}) = \mathfrak{C}^0_{\pm 4}\,, && \mathrm{gr}_{\pm 3}(\mathfrak{C}) = \mathfrak{C}^0_{\pm 3}\,, && \mathrm{gr}_{\pm 2}(\mathfrak{C}) = \mathfrak{C}^0_{\pm 2} \oplus \mathfrak{C}^1_{\pm 2} \oplus \mathfrak{C}^2_{\pm 2}\,, \\ &\mathrm{gr}_{\pm 1}(\mathfrak{C}) = \mathfrak{C}^0_{\pm 1} \oplus \mathfrak{C}^1_{\pm 1} \oplus \mathfrak{C}^2_{\pm 1}\,, && \mathrm{gr}_0(\mathfrak{C}) = \mathfrak{C}^0_0 \oplus \mathfrak{C}^1_0 \oplus \mathfrak{C}^2_0 \oplus \mathfrak{C}^3_0 \oplus \mathfrak{C}^4_0\,. \end{aligned}$$

where the $\mathfrak{C}^j_i$ *are given in Table 5 of appendix D. Further,*

$$\mathfrak{C}^j_i = \{C_{abcd} \in \mathfrak{C}^i : {}^{\mathfrak{C}}\Pi^k_i(C) = 0 \, \textit{for all } k \neq j\}/\mathfrak{C}^{i+1}\,.$$

for $|i| \leq 3$*, where the maps* ${}^{\mathfrak{C}}\Pi^j_i$ *are defined in appendix A.*

Finally, the $\mathfrak{p}$*-orbit structure of this decomposition can be expressed by means of the graph*

$$\begin{tikzcd}
 & & & & \mathfrak{C}^4_0 \arrow[dr] & & & & \\
 & & \mathfrak{C}^2_2 \arrow[r] \arrow[ddr] & \mathfrak{C}^2_1 \arrow[ur] \arrow[dr] \arrow[dddr] & & \mathfrak{C}^2_{-1} \arrow[r] \arrow[ddr] & \mathfrak{C}^2_{-2} \arrow[ddr] & & \\
 & & & & \mathfrak{C}^3_0 \arrow[ur] \arrow[dr] & & & & \\
\mathfrak{C}^0_4 \arrow[r] & \mathfrak{C}^0_3 \arrow[uur] \arrow[r] \arrow[ddr] & \mathfrak{C}^1_2 \arrow[uur] \arrow[ddr] & \mathfrak{C}^1_1 \arrow[ur] \arrow[dr] & & \mathfrak{C}^1_{-1} \arrow[uur] \arrow[ddr] & \mathfrak{C}^1_{-2} \arrow[r] & \mathfrak{C}^0_{-3} \arrow[r] & \mathfrak{C}^0_{-4} \\
 & & & & \mathfrak{C}^1_0 \oplus \mathfrak{C}^2_0 \arrow[uuur] \arrow[ur] \arrow[dr] & & & & \\
 & & \mathfrak{C}^0_2 \arrow[uur] \arrow[r] & \mathfrak{C}^0_1 \arrow[ur] \arrow[dr] & & \mathfrak{C}^0_{-1} \arrow[uur] \arrow[r] & \mathfrak{C}^0_{-2} \arrow[uur] & & \\
 & & & & \mathfrak{C}^0_0 \arrow[ur] & & & &
\end{tikzcd}$$

with the proviso that when $m = 2$*, the modules* $\mathfrak{C}^2_{\pm 2}$*,* $\mathfrak{C}^1_{\pm 1}$*,* $\mathfrak{C}^3_0$ *do not occur, and the isotopic modules* $\mathfrak{C}^1_0$ *and* $\mathfrak{C}^2_0$ *degenerate to the same module, and when* $m = 3$*, the module* $\mathfrak{C}^3_0$ *does not occur. Here, an arrow from* $\mathfrak{C}^j_i$ *to* $\mathfrak{C}^k_{i-1}$ *for some* i, j, k *implies that* $\breve{\mathfrak{C}}^j_i \subset \mathfrak{g}_1 \cdot \breve{\mathfrak{C}}^k_{i-1}$ *for any choice of irreducible* $\mathfrak{g}_0$*-modules* $\breve{\mathfrak{C}}^j_i$ *and* $\breve{\mathfrak{C}}^k_{i-1}$ *isomorphic to* $\mathfrak{C}^j_i$ *and* $\mathfrak{C}^k_{i-1}$ *respectively.*

4 The $\mathfrak{p}$-invariant classification of $\mathfrak{V} \otimes (\mathfrak{g}/\mathfrak{p})$

We can apply the above recipe starting from the $\mathfrak{p}$-module $\mathfrak{W} := \mathfrak{V} \otimes \mathfrak{g}/\mathfrak{p}$. Again, we assume $m > 2$, leaving the case $m = 2$ to appendix B. The $\mathfrak{p}$-invariant filtrations on $\mathfrak{V}$ and $\mathfrak{g}$ induces a $\mathfrak{p}$-invariant filtration on $\mathfrak{W}$

$$\mathfrak{W}^0 \subset \mathfrak{W}^{-1} \subset \mathfrak{W}^{-2} \subset \mathfrak{W}^{-3}\,,$$

where

$$\begin{aligned} &\mathfrak{W}^{-3} := \mathfrak{V}^{-1} \otimes \left(\mathfrak{g}^{-2}/\mathfrak{g}^0\right)\,, && \mathfrak{W}^{-2} := \left(\mathfrak{V}^{-1} \otimes \left(\mathfrak{g}^{-1}/\mathfrak{g}^0\right)\right) \oplus \left(\mathfrak{V}^0 \otimes \left(\mathfrak{g}^{-2}/\mathfrak{g}^0\right)\right)\,, \\ &\mathfrak{W}^{-1} := \left(\mathfrak{V}^0 \otimes \left(\mathfrak{g}^{-1}/\mathfrak{g}^0\right)\right) \oplus \left(\mathfrak{V}^1 \otimes \left(\mathfrak{g}^{-2}/\mathfrak{g}^0\right)\right)\,, && \mathfrak{W}^0 := \mathfrak{V}^1 \otimes \left(\mathfrak{g}^{-1}/\mathfrak{g}^0\right)\,. \end{aligned}$$

We can express an element of $\mathfrak{W}$ as $\Gamma_{abc}\xi^{bB}\xi^{cC}$ or $\Gamma_{abc}\xi^{bcB}$ (mod $\alpha_a\xi^B$). To assist the analysis, it is convenient to work in the splitting (2.14), and use the spinors ξ^A and η_A satisfying $\xi^A\eta_A = -\frac{1}{2}$, and

$u_a = 2\,\xi^{aA}\eta_A$ as in section 2. We then have

$$
\begin{aligned}
\Gamma_{abc}\xi^{bB}\xi^{cC} &= \xi_a{}^A\Gamma_A{}^{BC} - u_a\Gamma^{BC} + \eta_{aA}\Gamma^{ABC} + 2\xi_a{}^A\Gamma_{A:}{}^{[B}\xi^{C]} - 2u_a\Gamma^{[B}\xi^{C]} + 2\eta_{aA}\Gamma^{A:[B}\xi^{C]}\,, \\
\Gamma_{abc}\xi^{bcD} &= \left(\xi_a{}^A\Gamma_A{}^{EC} - u_a\Gamma^{EC} + \eta_{aA}\Gamma^{AEC}\right)\eta_{cC}\gamma^c{}_E{}^D - 2\left(\xi_a{}^A\Gamma_{A:}{}^D - u_a\Gamma^D + \eta_{aA}\Gamma^{A:D}\right) \\
&\qquad + 2\left(\xi_a{}^A\Gamma_{AB}{}^B - u_a\Gamma_B{}^B + \eta_{aA}\Gamma^A{}_B{}^B\right)\xi^D\,, \\
\gamma^a{}_D{}^A\Gamma_{abc}\xi^{bcD} &= 4\Gamma_{C:}{}^C\xi^A + 4\Gamma_C{}^{CA} - 2\Gamma^A + \eta_{aB}\gamma^a{}_C{}^A\left(\Gamma^{BC} - \Gamma^{B:C} + \Gamma^{C:B}\right) + \eta_{bB}\eta_{cC}\Gamma^{DBC}\gamma^{bc}{}_D{}^A \\
&\qquad - 2\Gamma_B{}^B\xi^A + 2\Gamma^A{}_B{}^B\,,
\end{aligned}
\tag{4.1}
$$

where $\Gamma_A{}^{BC} = \Gamma_A{}^{[BC]}$, $\Gamma^{ABC} = \Gamma^{A[BC]}$, $\Gamma_{AB}{}^C$, $\Gamma^A{}_B{}^C$, Γ^A, $\Gamma_{A:}{}^B$, $\Gamma_A{}^B$, $\Gamma^{A:B}$, $\Gamma^{AB} = \Gamma^{[AB]}$ are all elements of $\mathfrak{S}_{\frac{m-2}{2}}$ and $\mathfrak{S}_{-\frac{m-2}{2}}$ and tensor products thereof.

Remark 4.1 The colon : in $\Gamma_{A:}{}^B$ and $\Gamma^{A:B}$ does *not* denote any differential operation, but is simply a marker separating the '1-form' index from the 'Lie algebra' indices.

The module $\mathfrak{W}$ decomposes as a direct sum of $\mathfrak{g}_0$-modules $\breve{\mathfrak{W}}_i^j$, each of which determined by the components

$$
\begin{array}{lll}
\Gamma^{[ABC]} \in \breve{\mathfrak{W}}^0_{-3}\,, & \Gamma^{(AB)C} \in \breve{\mathfrak{W}}^1_{-3}\,, & \\
\Gamma^{AB} \in \breve{\mathfrak{W}}^0_{-2}\,, & \Gamma^{[A:B]} \in \breve{\mathfrak{W}}^1_{-2}\,, & \Gamma^{(A:B)} \in \breve{\mathfrak{W}}^2_{-2}\,, \\
\Gamma_B{}^{BA} \in \breve{\mathfrak{W}}^0_{-1}\,, & \Gamma_A{}^{BC} - \dfrac{2}{m-1}I_A^{[B|}\Gamma_D{}^{D|C]} \in \breve{\mathfrak{W}}^1_{-1}\,, & \Gamma^A \in \breve{\mathfrak{W}}^2_{-1}\,, \\
\Gamma_{A:}{}^A \in \breve{\mathfrak{W}}^0_0\,, & \Gamma_{A:}{}^B - \dfrac{1}{m}I_A^B\Gamma_{C:}{}^C \in \breve{\mathfrak{W}}^1_0\,. &
\end{array}
$$

To make the analysis invariant, we define

$$
\begin{aligned}
{}^{\mathfrak{W}}\Pi^0_{-3}(\Gamma) &:= \Gamma_{abc}\xi^{a[A}\xi^{bB}\xi^{cC}\xi^{D]}\,, \\
{}^{\mathfrak{W}}\Pi^1_{-3}(\Gamma) &:= \Gamma_{abc}\xi^{[A}\xi^{aB]}\xi^{b[C}\xi^{D]}\xi^{c[E}\xi^{F]} + \Gamma_{abc}\xi^{[C}\xi^{aD]}\xi^{b[A}\xi^{B]}\xi^{c[E}\xi^{F]}\,, \\
{}^{\mathfrak{W}}\Pi^0_{-2}(\Gamma) &:= \Gamma_{abc}\xi^{aA}\xi^{b[B}\xi^{cC}\xi^{D]}\,, \\
{}^{\mathfrak{W}}\Pi^1_{-2}(\Gamma) &:= \Gamma_{abc}\xi^{a[A}\xi^{bcB}\xi^{C]}\,, \\
{}^{\mathfrak{W}}\Pi^2_{-2}(\Gamma) &:= \Gamma_{abc}\xi^{[A}\xi^{aB]}\xi^{b[C}\xi^{D]}\xi^{cE} + \Gamma_{abc}\xi^{[C}\xi^{aD]}\xi^{b[A}\xi^{B]}\xi^{cE}\,, \\
{}^{\mathfrak{W}}\Pi^0_{-1}(\Gamma) &:= \begin{cases} \gamma^a{}_C{}^{[A|}\Gamma_{abc}\xi^{bcC}\xi^{|B]} - \xi^{a[A}\Gamma_{abc}\xi^{bcB]}\,, & m > 2\,, \\ \gamma^a{}_C{}^{[A|}\Gamma_{abc}\xi^{bcC}\xi^{|B]} - \xi^{a[A}\Gamma_{abc}\xi^{bcB]} + \tfrac{1}{2}\varepsilon^{AB}\xi^a{}_C\Gamma_{abc}\xi^{bcC}\,, & m = 2\,, \end{cases} \\
{}^{\mathfrak{W}}\Pi^1_{-1}(\Gamma) &:= \Gamma_{abc}\xi^{b[B}\xi^{cC}\xi^{D]} - \frac{1}{2(m-1)}\left(\gamma^d{}_A{}^{[B|}\Gamma_{dbc}\xi^{bcA}\xi^{|C} - \xi^{d[B}\Gamma_{dbc}\xi^{bcC}\right)\xi_a{}^{D]}\,, \\
{}^{\mathfrak{W}}\Pi^2_{-1}(\Gamma) &:= \Gamma_{abc}\xi^{aA}\xi^{bc[B}\xi^{C]}\,, \\
{}^{\mathfrak{W}}\Pi^0_0(\Gamma) &:= \gamma^a{}_C{}^A\Gamma_{abc}\xi^{bcC}\xi^B - \xi^{aA}\Gamma_{abc}\xi^{bcB}\,, \\
{}^{\mathfrak{W}}\Pi^1_0(\Gamma) &:= \Gamma_{abc}\xi^{bc[B}\xi^{C]} + \frac{1}{2m}\left(\gamma^d{}_A{}^{[B|}\Gamma_{dbc}\xi^{bcA}\xi_a{}^{|C]} - \xi^{d[B|}\Gamma_{dbc}\xi^{bcA}\gamma_{aA}{}^{|C]}\right)\,,
\end{aligned}
$$

where $\Gamma_{abc} \in \mathfrak{V} \otimes \mathfrak{g}$. We have been careful to distinguish the cases $m > 2$ and $m = 2$. In the latter case, we have made use of the $\mathrm{Spin}(5,\mathbb{C})$-invariant skewsymmetric bilinear forms ε_{AB} and ε^{AB}.

Proposition 4.2 *The associated graded* $\mathfrak{p}$*-module* $\mathrm{gr}(\mathfrak{W})$ *of* $\mathfrak{W}$ *decomposes into a direct sum*

$$
\begin{array}{ll}
\mathrm{gr}_{-3}(\mathfrak{W}) \cong \mathfrak{W}^0_{-3} \oplus \mathfrak{W}^1_{-3}\,, & \mathrm{gr}_{-2}(\mathfrak{W}) \cong \mathfrak{W}^0_{-2} \oplus \mathfrak{W}^1_{-2} \oplus \mathfrak{W}^2_{-2}\,, \\
\mathrm{gr}_{-1}(\mathfrak{W}) \cong \mathfrak{W}^0_{-1} \oplus \mathfrak{W}^1_{-1} \oplus \mathfrak{W}^2_{-1}\,, & \mathfrak{W}^0 \cong \mathfrak{W}^0_0 \oplus \mathfrak{W}^1_0\,,
\end{array}
$$

of irreducible $\mathfrak{p}$*-modules, where the* $\mathfrak{W}^j_i$ *are given in Table 6 of appendix D. Further,*

$$\mathfrak{W}^j_i = \left\{ \Gamma \in \mathfrak{W}^i : {}^{\mathfrak{W}}\Pi^k_i(\Gamma) = 0 \,, \textit{for all } k \neq j \right\} / \mathfrak{W}^{i+1} \,, \qquad\qquad i = -3, -2, -1, 0 \,.$$

Finally, the $\mathfrak{p}$*-orbit structure of this decomposition can be expressed by means of the graph*

$$\begin{tikzcd}
 & \mathfrak{W}^2{}_{-1} \arrow[r] \arrow[ddr] \arrow[ddddr] & \mathfrak{W}^2{}_{-2} \arrow[dr] & \\
\mathfrak{W}^1_0 \arrow[ur] \arrow[dr] \arrow[dddr] & & & \mathfrak{W}^1{}_{-3} \\
 & \mathfrak{W}^1{}_{-1} \arrow[r] \arrow[ddr] & \mathfrak{W}^1{}_{-2} \arrow[ur] \arrow[dr] & \\
\mathfrak{W}^0_0 \arrow[uuur] \arrow[dr] & & & \mathfrak{W}^0{}_{-3} \\
 & \mathfrak{W}^0{}_{-1} \arrow[r] \arrow[uur, dotted] & \mathfrak{W}^0{}_{-2} \arrow[uuur] \arrow[ur] &
\end{tikzcd}$$

with the proviso that $\mathfrak{W}^1{}_{-1}$ *and the dotted arrow only occur when* $m > 2$*. Here, an arrow from* $\mathfrak{W}^j_i$ *to* $\mathfrak{W}^k_{i-1}$ *for some* i, j, k *implies that* $\breve{\mathfrak{W}}^j_i \subset \mathfrak{g}_1 \cdot \breve{\mathfrak{W}}^k_{i-1}$ *for any choice of irreducible* $\mathfrak{g}_0$*-modules* $\breve{\mathfrak{W}}^j_i$ *and* $\breve{\mathfrak{W}}^k_{i-1}$ *isomorphic to* $\mathfrak{W}^j_i$ *and* $\mathfrak{W}^k_{i-1}$ *respectively.*

5 Differential geometry of pure spinor fields

Throughout, $(\mathcal{M}, g)$ will denote an n-dimensional complex Riemannian manifold, where $n = 2m+1$, i.e. a complex manifold equipped with a holomorphic non-degenerate symmetric bilinear form g_{ab}. The holomorphic tangent and cotangent bundles will be denoted by $\mathrm{T}\mathcal{M}$ and $\mathrm{T}^*\mathcal{M}$ respectively. Thus, g_{ab} is a holomorphic section of $\odot^2\mathrm{T}^*\mathcal{M}$. The structure group of the frame bundle $\mathcal{FM}$ of $\mathcal{M}$ is reduced to $\mathrm{SO}(2m+1, \mathbb{C})$.

The complex Riemannian manifold $(\mathcal{M}, g)$ admits a unique torsion-free connection preserving g_{ab}, the holomorphic Levi-Civita connection, which, together with its associated covariant derivative, will be denoted ∇_a. For convenience, we can choose an orthonormal frame $\{\delta^a_{\mathbf{a}}\}$ with dual $\{\delta^{\mathbf{a}}_a\}$, where bold lower case Roman indices run from 1 to n. The connection ∇_a can then be represented by the $\mathfrak{g}$-valued 1-form $\Gamma_{\mathbf{ab}}{}^{\mathbf{c}} := (\delta^a_{\mathbf{a}} \nabla_a \delta^b_{\mathbf{b}}) \delta^{\mathbf{c}}_b$. Choosing a connnection ∂_a compatible with g_{ab}, $\{\delta^a_{\mathbf{a}}\}$ and $\{\delta^{\mathbf{a}}_a\}$, the connections ∇_a and ∂_a are then related via

$$\nabla_a V^b = \partial_a V^b + \Gamma_{ac}{}^b V^c \,, \tag{5.1}$$

for any holomorphic vector field V^a, and where $\Gamma_{ab}{}^c := \delta^{\mathbf{a}}_a \delta^{\mathbf{b}}_b \Gamma_{\mathbf{ab}}{}^{\mathbf{c}} \delta^c_{\mathbf{c}}$. The Riemann curvature R_{abcd} of ∇_a is defined by

$$2 \nabla_{[a} \nabla_{b]} V^c = R_{abd}{}^c V^d \,,$$

for any holomorphic vector field V^a, and similarly for holomorphic sections of $\mathrm{T}^*\mathcal{M}$, and tensor products. The Riemann tensor splits into irreducible $\mathrm{SO}(2m+1, \mathbb{C})$-invariant parts,

$$R_{abcd} = C_{abcd} + \frac{4}{n-2} \Phi_{[a|[c} g_{d]|b]} + \frac{2}{n(n-1)} R g_{a[c} g_{d]b} \,, \tag{5.2}$$

where C_{abcd} is the Weyl tensor, Φ_{ab} is the tracefree part of the Ricci tensor $R_{ab} := R_{acb}{}^{c}$, and $R := R_{a}{}^{a}$ is the Ricci scalar.

We shall assume that $(\mathcal{M}, g)$ is equipped with a spin structure – see [LM89] for definitions and topological restrictions. The spinor bundle on $\mathcal{M}$ admits a canonical connection as a lift of the Levi-Civita connection, and for this reason, this spinor connection will also be denoted ∇_a. It preserves the Clifford module of the spinor bundle in the sense that

$$\nabla_a \left(V^b \gamma_{bC}{}^{B} \xi^C \right) = \left(\nabla_a V^b \right) \gamma_{bC}{}^{A} \xi^C + V^b \gamma_{bC}{}^{A} \nabla_a \xi^C \, ,$$

for any holomorphic vector field V^a and holomorphic spinor field ξ^A, and similarly for dual spinor fields. The relation between the lifts of ∂_a introduced in (5.1) and ∇_a to the spinor bundle is given by

$$\nabla_a \xi^A = \partial_a \xi^A - \frac{1}{4} \Gamma_{abc} \gamma^{bc}{}_{B}{}^{A} \xi^B \, . \tag{5.3}$$

Similarly, the curvature tensor of the spinor bundle can be expressed as a lift of the Riemann curvature tensor, i.e.

$$2 \nabla_{[a} \nabla_{b]} \xi^A = - \frac{1}{4} R_{abcd} \gamma^{cd}{}_{B}{}^{A} \xi^B \, .$$

for any holomorphic spinor field ξ^A, and similarly for dual spinor fields.

5.1 Projective pure spinor fields

A spinor field ξ^A is said to be pure if at every point $p \in \mathfrak{M}$, ξ^A defines a totally null m-dimensional vector subspace

$$\mathcal{N}_p := \{ X^a \in \mathrm{T}_p \mathcal{M} : X^a \xi^{aA} = 0 \} \, ,$$

of $\mathrm{T}_p\mathcal{M}$. Here, $\xi^{aA} := \xi^B \gamma^a{}_B{}^A$ as before. Thus, a pure spinor field defines a γ-plane distribution, and conversely, any such distribution determines a pure spinor field up to scale. This motivates the following definition, which, it is worth pointing out, does not require the existence of a spin structure on $\mathcal{M}$.

Definition 5.1 The *projective pure spinor bundle* $\mathrm{Gr}_m(\mathrm{T}\mathcal{M}, g)$ is the bundle over $\mathcal{M}$ whose fiber over a point p of $\mathcal{M}$ is isomorphic to the $\frac{1}{2}m(m+1)$-dimensional family $\mathrm{Gr}_m(\mathrm{T}_p\mathcal{M}, g)$ of γ-planes in $\mathrm{T}_p\mathcal{M}$.

From now on, we assume the existence of a holomorphic pure spinor field ξ^A on $\mathcal{M}$, i.e. a (global) holomorphic section of $\mathrm{Gr}_m(\mathcal{M}, g)$, with associated holomorphic distribution $\mathcal{N}$. This reduces the structure group of the frame bundle to P, the parabolic Lie subgroup of G stabilising $[\xi^A]$, as described in section 2. It is a standard result, e.g. [Sal89, ČS09b] that one can construct holomorphic vector bundles from the representation of P or its Lie algebra $\mathfrak{p}$. Thus, if $\mathfrak{C}$ is the irreducible G-module of tensors with Weyl symmetries, its associated vector bundle $\mathcal{C} := \mathcal{F}\mathcal{M} \times_G \mathfrak{C}$ admits a filtration of P-invariant subbundles $\mathcal{C}^i := \mathcal{F}\mathcal{M} \times_P \mathfrak{C}^i$, and similarly, to the associated graded $\mathfrak{p}$-modules $\mathrm{gr}(\mathfrak{C})$, $\mathfrak{g}_0$-modules $\mathfrak{C}_i$, and their irreducibles correspond associated graded vector bundles, and so on.

Remark 5.2 For convenience, we can choose a spinor field η_A dual to ξ^A as described in section 2, and let $\{\delta^{a\mathbf{A}}\}$ and $\{\delta^a_{\mathbf{A}}\}$ be bases of holomorphic vector fields annihilating ξ^A and η_A respectively, and dual to each other, i.e. $\delta^a_{\mathbf{A}} \delta^{\mathbf{B}}_a = \delta^{\mathbf{B}}_{\mathbf{A}}$. Here, bold upper cases Roman indices run from 1 to m. Defining $u_a := 2\xi^{aA}\eta_A$, together with a null frame $\{\delta^a_{\mathbf{A}}, \delta^{a\mathbf{A}}, u_a\}$, then the tensors Γ^{ABC}, $\Gamma^{A:B}$, etc... appearing in (4.1) can be thought of as the abstract versions of the components $\Gamma^{\mathbf{ABC}} = \left(\delta^{a\mathbf{A}} \nabla_a \delta^{b\mathbf{B}} \right) \delta^{\mathbf{C}}_b$, $\Gamma_{\mathbf{A}:}{}^{\mathbf{B}} = \left(\delta^a_{\mathbf{A}} \nabla_a \delta^{b\mathbf{B}} \right) u_b$, and so on, of the connection 1-form $\Gamma_{ab}{}^{c}$.

Now, choosing our connection ∂_a to preserve ξ^A and η_A, the covariant derivative (5.3).of ξ^A reduces to

$$\nabla_a \xi^D = - \frac{1}{4} \Gamma_{abc} \xi^A \gamma^{bc}{}_{A}{}^{D} \, .$$

Remark 5.3 This section applies equally to a $(2m+1)$-dimensional smooth manifold equipped with a smooth metric of split signature. In this case, the projective pure spinor field $[\xi^A]$ is assumed to be *real* in the sense that its associated null m-plane distribution is real.

5.1.1 The intrinsic torsion of a projective pure spinor field

The *intrinsic torsion* [Sal89] of the projective pure spinor field $[\xi^A]$ measures the failure of $[\xi^A]$ to be parallel with respect to the Levi-Civita connection, i.e.

$$\nabla_a[\xi^A] = 0 \,, \qquad \text{i.e.} \qquad \nabla_a \xi^A = \alpha_a \xi^A \,, \tag{5.4}$$

for some 1-form α_a, or equivalently, the failure of the connection 1-form taking values in $\mathfrak{p}$. Since ∇_a takes values in $\mathfrak{g}$, this measure is precisely obtained by consideration of the module $\mathfrak{W} = \mathfrak{V} \otimes (\mathfrak{g}/\mathfrak{p})$, which we defined in section 4. The tensor of interest will be given by

$$\Gamma_{abc} \xi^{bB} \xi^{cC} \qquad \in \qquad \mathfrak{V}^{-1} \otimes \wedge^2 \mathfrak{S}^{-\frac{m-2}{2}} \cong \mathfrak{W} \,.$$

We shall then be able to apply Proposition 4.2 and the maps ${}^{\mathfrak{W}}\Pi_i^j$ describing the various irreducibles $\mathfrak{p}$-modules of $\mathfrak{W}$ to characterise the intrinsic torsion of $[\xi^A]$. Before that, in order to give covariant expressions for these, we compute, from (5.3) and (2.18), the formula

$$\left(\nabla_a \xi^{bB}\right) \xi_b{}^C = -\left(\nabla_a \xi^B\right) \xi^C + \Gamma_{abc} \xi^{bB} \xi^{cC}$$

from which we deduce

$$\begin{aligned}
\left(\nabla_a \xi^{b(B}\right) \xi_b{}^{C)} &= -\left(\nabla_a \xi^{(B}\right) \xi^{C)} \,, & \left(\nabla_a \xi^{b[B}\right) \xi_b{}^{C]} &= -\left(\nabla_a \xi^{[B}\right) \xi^{C]} + \Gamma_{abc} \xi^{bB} \xi^{cC} \,, \\
\left(\nabla_a \xi^{b[B}\right) \xi_b{}^C \xi^{D]} &= \Gamma_{abc} \xi^{b[B} \xi^{cC} \xi^{D]} \,, & \left(\nabla_a \xi^{b[A}\right) \xi^{B]} \xi_b{}^{[C} \xi^{D]} &= \Gamma_{abc} \xi^{b[A} \xi^{B]} \xi^{c[C} \xi^{D]} \,.
\end{aligned}$$

The first of these identities is trivially satisfied by virtue of the purity condition.

Proposition 5.4 *Pointwise, the intrinsic torsion of* $[\xi^A]$ *projects trivially into*

- $\mathfrak{W}^0_{-3}$ *if and only if*

$$\left(\xi^{a[A} \nabla_a \xi^{bB}\right) \xi_b{}^C \xi^{D]} = 0 \,; \tag{5.5}$$

- $\mathfrak{W}^1_{-3}$ *if and only if*

$$\xi^{[A} \left(\xi^{aB]} \nabla_a \xi^{b[C}\right) \xi^{D]} \xi_b{}^{[E} \xi^{F]} + \xi^{[C} \left(\xi^{aD]} \nabla_a \xi^{b[A}\right) \xi^{B]} \xi_b{}^{[E} \xi^{F]} = 0 \,; \tag{5.6}$$

- $\mathfrak{W}^0_{-2}$ *and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-2}$ *if and only if*

$$\left(\xi^{aA} \nabla_a \xi^{b[B}\right) \xi_b{}^C \xi^{D]} = 0 \,; \tag{5.7}$$

- $\mathfrak{W}^1_{-2}$ *and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-2}$ *if and only if*

$$\left(\xi^{a[A} \nabla_a \xi^B\right) \xi^{C]} = 0 \,; \tag{5.8}$$

- $\mathfrak{W}^2_{-2}$ *and* $\mathfrak{W}^1_{-3}$ *if and only if*

$$\xi^{[A} \left(\xi^{aB]} \nabla_a \xi_b{}^E\right) \xi^{b[C} \xi^{D]} + \xi^{[C} \left(\xi^{aD]} \nabla_a \xi_b{}^E\right) \xi^{b[A} \xi^{B]} = 0 \,; \tag{5.9}$$

- $\mathfrak{W}^0{}_{-1}$, $\mathfrak{W}^0{}_{-2}$, $\mathfrak{W}^1{}_{-2}$ *and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-2}$ *if and only if*

$$\left(\nabla_a \xi^{a[A}\right) \xi^{B]} - \left(\xi^{a[A} \nabla_a \xi^{B]}\right) = 0\,; \tag{5.10}$$

- $\mathfrak{W}^1{}_{-1}$, $\mathfrak{W}^0{}_{-2}$, $\mathfrak{W}^1{}_{-2}$ *and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-2}$ *if and only if*

$$\left(\nabla_a \xi^{b[B}\right) \xi_b{}^C \xi^{D]} + \frac{2}{m-1}\left(\left(\nabla_b \xi^{b[B}\right) \xi^C - \xi^{b[B} \nabla_b \xi^C\right) \xi_a{}^{D]} = 0\,; \tag{5.11}$$

- $\mathfrak{W}^2{}_{-1}$ *and* $\mathfrak{W}^{-2}/\mathfrak{W}^{-1}$ *if and only if*

$$\left(\xi^{aA} \nabla_a \xi^{[B}\right) \xi^{C]} = 0\,; \tag{5.12}$$

- $\mathfrak{W}^0_0$, $\mathfrak{W}^0{}_{-1}$, $\mathfrak{W}^2{}_{-1}$ *and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-1}$ *if and only if*

$$\left(\nabla_a \xi^{aA}\right) \xi^B - \xi^{aA} \nabla_a \xi^B = 0\,; \tag{5.13}$$

- $\mathfrak{W}^1_0$ *and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-0}$ *if and only if*

$$\left(\nabla_a \xi^{[B}\right) \xi^{C]} - \frac{2}{m}\left(\left(\nabla_b \xi^{b[B}\right) \xi_a{}^{C]} - \xi^{b[B} \nabla_b \xi_a{}^{C]}\right) = 0\,. \tag{5.14}$$

Further, when $m = 2$*, the intrinsic torsion of* $[\xi^A]$ *projects trivially into* $\mathfrak{W}^0{}_{-1}$*,* $\mathfrak{W}^0{}_{-2}$*, and* $\mathfrak{W}^{-3}/\mathfrak{W}^{-2}$ *if and only if*

$$\left(\nabla_a \xi^{a[A}\right) \xi^{B]} - \xi^{a[A} \nabla_a \xi^{B]} + \frac{1}{2} \varepsilon^{AB} \xi^a{}_C \nabla_a \xi^C = 0\,, \tag{5.15}$$

where $\varepsilon^{AB} = \varepsilon^{[AB]}$ *is the spin invariant bilinear form on* $\mathfrak{S}^*$*.*

5.1.2 Geometric properties

We come to one of the main motivations of this article, finding differential conditions on $[\xi^A]$ to describe the integrability properties of its associated γ-plane distribution $\mathcal{N}$. Denote by $\mathcal{N}^\perp$ the orthogonal complement of $\mathcal{N}$.

Proposition 5.5 *Let* $[\xi^A]$ *be a projective pure spinor field with associated* γ*-plane distribution* $\mathcal{N}$ *as before. Then*

- $[\Gamma(\mathcal{N}), \Gamma(\mathcal{N})] \subset \Gamma(\mathcal{N}^\perp)$*, i.e. if and only if*

$$\xi^{[A}\left(\xi^{aB]} \nabla_a \xi^{b[C}\right) \xi_b{}^D \xi^{E]} = 0\,; \tag{5.16}$$

- $[\Gamma(\mathcal{N}), \Gamma(\mathcal{N})] \subset \Gamma(\mathcal{N})$ *if and only if*

$$\left(\xi^{a[A} \nabla_a \xi^B\right) \xi^{C]} = 0\,; \tag{5.8}$$

- $[\Gamma(\mathcal{N}^\perp), \Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N}^\perp)$ *if and only if*

$$\left(\xi^{a[A} \nabla_a \xi^{bB]}\right) \xi_b{}^{[C} \xi^{D]} = 0\,. \tag{5.17}$$

- $[\Gamma(\mathcal{N}),\Gamma(\mathcal{N})] \subset \Gamma(\mathcal{N})$ *and* $[\Gamma(\mathcal{N}^\perp),\Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N}^\perp)$ *if and only if*

$$\left(\xi^{aA}\nabla_a \xi^{b[B}\right)\xi_b{}^C\xi^{D]} = 0\,; \qquad\qquad \xi^{[A}\left(\xi^{aB]}\nabla_a \xi^{[C}\right)\xi^{D]} = 0\,; \tag{5.18}$$

- $[\Gamma(\mathcal{N}^\perp),\Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N})$ *if and only if*

$$\left(\xi^{aA}\nabla_a \xi^{[B}\right)\xi^{C]} = 0\,. \tag{5.12}$$

Remark 5.6 Conditions (5.16) and (5.18) are equivalent to the intrinsic torsion of the P-structure lying in the module $\mathfrak{W}^{-2}$ and $\mathfrak{W}^{-1}$ respectively. Further, condition (5.17) implies that the intrinsic torsion belongs to the module $\mathfrak{W}^{-2}$ and that it projects trivially into $\mathfrak{W}^2_{-2}$.

Proof. One can apply the Frobenius theorem on vector fields of the form $V^a = \xi^{aA}v_A$, where v_A lies in $\mathfrak{S}^{-\frac{m}{2}}/\mathfrak{S}^{-\frac{m-4}{2}}$ or $\mathfrak{S}^{-\frac{m-2}{2}}/\mathfrak{S}^{-\frac{m-4}{2}}$, and, with reference to (4.1), we note that each of the above conditions is equivalent to

- $\Gamma^{ABC} = 0$,
- $\Gamma^{ABC} = \Gamma^{[A:B]} = 0$,
- $\Gamma^{ABC} = 0$ and $\Gamma^{A:B} = \Gamma^{AB}$, (in particular, $\Gamma^{(A:B)} = 0$),
- $\Gamma^{ABC} = \Gamma^{AB} = \Gamma^{A:B} = 0$,
- $\Gamma^{ABC} = \Gamma^{AB} = \Gamma^{A:B} = \Gamma^{A} = 0$,

respectively. □

Definition 5.7 We say that ξ^A is

- *almost foliating* if $[\Gamma(\mathcal{N}),\Gamma(\mathcal{N})] \subset \Gamma(\mathcal{N}^\perp)$;
- *γ-foliating* if $[\Gamma(\mathcal{N}),\Gamma(\mathcal{N})] \subset \Gamma(\mathcal{N})$;
- *$\gamma^\perp$-foliating* if $[\Gamma(\mathcal{N}^\perp),\Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N}^\perp)$;
- *foliating* if it is both γ-foliating and $\gamma^\perp$-foliating;
- *strongly foliating* if $[\Gamma(\mathcal{N}^\perp),\Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N})$.

Conformal invariance With reference to appendix C, we prove

Proposition 5.8 *Conditions* (5.5)*,* (5.6)*,* (5.7)*,* (5.8)*,* (5.9)*,* (5.11) *and* (5.17)*, (and thus* (5.16)*,* (5.18)*) are conformally invariant. Suppose further that ξ^A satisfies* (5.11) *and*

$$\left(\nabla_a \xi^{a[A}\right)\xi^{B]} - \xi^{a[A}\nabla_a \xi^{B]} = (m-1)\xi^{[A}\xi^{aB]}\nabla_a f\,,$$

for some complex analytic function f. Then $(\mathcal{M}, g)$ is conformal to a complex Riemannian manifold admitting a parallel γ-plane distribution, i.e.

$$\left(\nabla_a \xi^{b[B}\right)\xi_b{}^C\xi^{D]} = 0\,. \tag{5.19}$$

Remark 5.9 The second part of the above proposition should be compared with its even-dimensional counterpart given in [TC12b], which was argued to be a complex Riemannian analogue of Hermitian manifolds locally conformal to Kähler [FFS94], in the sense of 'conformal to a complex Riemmannian manifold admitting a recurrent pure spinor field'. Proposition 5.8 shows that in odd dimensions, the appropriate generalisation is (5.19), not (5.4).

Curvature conditions The integrability conditions for these equations can easily be computed by differentiation a second time and commuting the covariant derivatives.

Proposition 5.10 *Suppose that ξ^A is foliating. Then*

$$\xi^{[A} \xi^{aB} \xi^{bC]} \xi^{cD} \xi^{dE} C_{abcd} = 0 \, .$$

If ξ^A is strongly foliating, then

$$\xi^{aA} \xi^{bB} \xi^{cC} \xi^{dD} R_{abcd} = 0 \, , \qquad \xi^{[A} \xi^{aB} \xi^{bC]} \xi^{cD} \xi^{dE} C_{abcd} = 0 \, ,$$

and

$$\xi^{[A} \xi^{aB]} \Phi_{ab} \xi^{b[C} \xi^{D]} = 0 \qquad \Longleftrightarrow \qquad \xi^{aA} \xi^{bB} \xi^{cC} \xi^{dD} C_{abcd} = 0 \, .$$

Proposition 5.11 *Suppose that ξ^A satisfies*

$$\left(\nabla_a \xi^{b[B} \right) \xi_b{}^C \xi^{D]} = 0 \, , \tag{5.19}$$

i.e. the distribution $\mathcal{N}$ is parallel with respect to the Levi-Civita connection. Then

$$\xi^{[A} \xi^{aB} \xi^{bC]} R_{abcd} \xi^{cD} = 0 \, , \tag{5.20}$$

$$\xi^{[A} \xi^{aB} \xi^{bC]} C_{abcd} \xi^{cdD} = 0 \, . \tag{5.21}$$

Proof. We rewrite equation (5.19) as

$$\nabla_a \xi^A = \beta_a{}^A + \alpha_a \xi^A \, ,$$

for some 1-form α_a and $\mathfrak{S}_{\frac{m-2}{2}}$-valued 1-form $\beta_a{}^A$ such that $\xi^{[A} \xi^{aB]} \beta_a{}^C = 0$. Taking a second covariant derivative, commuting them, and using the identity (2.18) yield

$$R_{abcd} \xi^{cC} \xi^{dD} = 8 \left(\nabla_{[a} \beta_{b]}{}^{[C} \right) \xi^{D]} + 8 \beta_{[a}{}^{[C} \beta_{b]}{}^{D]} + 8 \alpha_{[a} \beta_{b]}{}^{[C} \xi^{D]} \, .$$

from which we obtain (5.20). To obtain (5.21), we use the decomposition of the Riemann tensor. □

Proposition 5.12 *Suppose that ξ^A is recurrent. Then*

$$\xi^{aA} \xi^{bB} R_{abcd} = 0 \, , \tag{5.22}$$

$$\xi^{aA} \xi^{bB} R_{ab} = 0 \, , \tag{5.23}$$

$$\xi^{aA} \xi^{b[B} \xi^{C]} \Phi_{ab} = 0 \, , \tag{5.24}$$

$$\xi^{aA} \xi^{bB} \xi^{c[C} \xi^{D]} C_{abcd} = 0 \, . \tag{5.25}$$

and in addition, if $m \geq 3$,

$$^{\mathfrak{C}}\Pi^3_0(C) = 0 \, .$$

Further, $\Pi^0_0(\Phi) = 0$ if and only if $R = 0$, and

$$\begin{aligned}
R = 0 \qquad & \Longleftrightarrow \qquad {}^{\mathfrak{C}}\Pi^0_0(C) = 0 \, , \\
{}^{\mathfrak{F}}\Pi^1_0(\Phi) = 0 \qquad & \Longleftrightarrow \qquad {}^{\mathfrak{C}}\Pi^2_0(C) = 0 \, , \\
{}^{\mathfrak{F}}\Pi^1_0(\Phi) = 0 \, , R = 0 \qquad & \Longleftrightarrow \qquad {}^{\mathfrak{C}}\Pi^0_0(C) = {}^{\mathfrak{C}}\Pi^1_0(C) = {}^{\mathfrak{C}}\Pi^2_0(C) = 0 \, , \\
{}^{\mathfrak{F}}\Pi^0_1(\Phi) = 0 \qquad & \Longleftrightarrow \qquad {}^{\mathfrak{C}}\Pi^0_1(C) = {}^{\mathfrak{C}}\Pi^1_1(C) = 0 \, .
\end{aligned}$$

Proof. Taking a second covariant derivative of (5.4), and commuting yield

$$-\frac{1}{4}R_{abcd}\xi^{cdA} = \left(\nabla_{[a}\,\alpha_{\,b]}\right)\xi^A \,,$$

which is equivalent to equation (5.22). Equations (5.23) and (5.24) are then obtained by contracting equation (5.22) with $\gamma^{cd}{}_A{}^B$, and using the expression relating R_{ab} and Φ_{ab} for which we find

$$\Phi_{ab}\xi^{aA}\xi^{bB} = \frac{1}{n}R\xi^A\xi^B \,.$$

Finally, using the decomposition of the Riemann tensor yields equation (5.25), and when $m \geq 3$,

$$\begin{aligned}
\xi^{cC}C_{c[ab]d}\xi^{dD} &= \frac{2}{n-2}\xi_{[a}{}^{[C}\Phi_{b]d}\xi^{dD]} + \frac{1}{n(n-1)}R\xi_{[a}{}^{C}\xi_{b]}{}^{D} \,, \\
C_{bcad}\xi^{bcB}\xi^{dD} &= -\frac{2(n-4)}{n-2}\xi^B\xi^{dD}\Phi_{ad} - \frac{2}{n-2}\xi^D\xi^{dB}\Phi_{ad} - \frac{2(n-2)}{n(n-1)}R\xi^B\xi_a{}^D + \frac{2}{n(n-1)(n-2)}R\xi^D\xi_a{}^B \,, \\
C_{bcad}\xi^{adA}\xi^{bcB} &= -2\left(\frac{n-3}{n-2}\right)R\xi^A\xi^B \,, \\
C_{abcd}\xi^{bB}\xi^{cC}\xi^{dD} &= -\frac{2}{n-2}\xi^B\xi^{[C}\Phi_{ad}\xi^{dD]} + \frac{2}{n(n-1)(n-2)}R\xi^B\xi^{[C}\xi_a{}^{D]} = -\frac{2}{n-2}\xi^{B\,\mathfrak{F}}\Pi^1_0(\Phi)_a{}^{CD} \,,
\end{aligned}$$

which, on referring to the maps ${}^{\mathfrak{C}}\Pi^j_i$, completes the proof. ☐

5.2 Spinorial differential equations

5.2.1 Parallel pure spinor fields

For reference, we recall the standard fact concerning parallel spinors – see for instance [Bry00].

Proposition 5.13 *Let ξ^A be a parallel spinor field, i.e. $\nabla_a\xi^A = 0$. Then*

$$\begin{aligned}
R_{abcd}\xi^{cdD} &= 0 \,, \\
R_{ab}\xi^{bB} = \mathrm{P}_{ab}\xi^{bB} = \Phi_{ab}\xi^{bB} &= 0 = R \,, \\
C_{abcd}\xi^{cdD} &= 0 \,.
\end{aligned}$$

The covariant derivative of a spinor field splits into two irreducible parts, which we shall presently turn our attention to.

5.2.2 Simple pure zero-rest-mass fields

The smaller irreducible part of the covariant derivative of a spinor field ξ^A leads to *(Weyl-)Dirac equation*

$$\gamma^a{}_A{}^B\nabla_a\xi^A = 0 \,, \tag{5.26}$$

In contrast to the even-dimensional, this equation admits not one, but *two* generalisations to irreducible spinor fields of higher valence.

Definition 5.14 Let $\phi^{A_1A_2\ldots A_k} = \phi^{(A_1A_2\ldots A_k)}$ be a holomorphic irreducible spinor field in the sense that

$$\gamma_{aA_1}{}^C\gamma_{aA_2}{}^D\phi^{A_1A_2A_3\ldots A_k} = -\phi^{CDA_3\ldots A_k} \,.$$

Then $\phi^{A_1\ldots A_k}$ is said to be a *zero-rest-mass (zrm) field*, respectively, *co-zero-rest-mass (co-zrm) field* if it satisfies

$$\gamma^a{}_B{}^{(A_1}\nabla_a\phi^{A_2\ldots\,A_k)B} = 0 \,, \tag{5.27}$$

respectively,

$$\gamma^{a}{}_{B}{}^{[A_1}\nabla_a\phi^{A_2]A_3\dots A_k B} = 0\,. \tag{5.28}$$

Remark 5.15 When $k = 2$, an irreducible spinor field as above is simply an m-form. Equation (5.27), respectively, (5.28) are then equivalent to this m-form to be closed, respecitvely, *co-closed*, hence the use of terminology.

Equations (5.27) and (5.28) are conformally invariant provided that $\phi^{A_1\dots A_k}$ is of conformal weight $-\frac{n+k-1}{2}$ in the former case, and $-\frac{n+2k-1}{2}$ in the latter case. In particular, a solution of both (5.27) and (5.28), i.e.

$$\gamma^{a}{}_{B}{}^{A_1}\nabla_a\phi^{A_2\dots A_k B} = 0\,, \tag{5.29}$$

is not conformally invariant. This equation in the case $k = 2$ corresponds to a closed and co-closed m-form.

The integrability condition on the existence of solutions of valence greater than two to equations (5.27) and (5.28) is given by the following lemma.

Proposition 5.16 *Assume $k > 2$. Let $\phi^{A_1A_2\dots A_k}$ be a solution of* (5.27) *or* (5.28). *Then*

$$\gamma^{a}{}_{C_1}{}^{A}\gamma^{b}{}_{C_2}{}^{B}C_{abcd}\gamma^{cd}{}_{D}{}^{(C_3}\phi^{C_4\dots C_k)C_1C_2D} = 0\,. \tag{5.30}$$

If $\phi^{A_1A_2\dots A_k}$ is a solution of (5.29), *then we have in addition*

$$\gamma^{b}{}_{C_2}{}^{[A|}\Phi_{bd}\gamma^{d}{}_{D}{}^{(C_3}\phi^{C_4\dots C_k)C_2D|B]} = 0\,. \tag{5.31}$$

Proof. Equations (5.27), (5.28) and (5.29) can be rewritten as

$$\gamma^{a}{}_{B}{}^{A_1}\nabla_a\phi^{A_2\dots A_k B} = \psi^{A_1A_2\dots A_k}\,,$$

where $\psi^{(A_1A_2\dots A_k)} = 0$, $\psi^{[A_1A_2]A_3\dots A_k} = 0$, and $\psi^{A_1A_2\dots A_k} = 0$ respectively. Taking a second covariant derivative and commuting lead to

$$\begin{aligned}(k-2)\,\gamma^{a}{}_{C_1}{}^{A}\gamma^{b}{}_{C_2}{}^{B}C_{abcd}\gamma^{cd}{}_{D}{}^{(C_3}\phi^{C_4\dots C_k)C_1C_2D} - 4(k-2)\,\gamma^{b}{}_{C_2}{}^{[A|}\mathrm{P}_{bd}\gamma^{d}{}_{D}{}^{(C_3}\phi^{C_4\dots C_k)C_2D|B]}& \\ = 2\,\gamma^{a}{}_{D}{}^{[A}\nabla_a\psi^{B]C_3C_4\dots C_kD}&\,.\end{aligned}$$

By the conformal invariance of (5.27) and (5.28), each term on the LHS must vanish identically, hence (5.30). When (5.29) holds, conformal invariance is broken, and one has the additional constraint (5.31). ☐

A spinor field $\phi^{A_1A_2\dots A_k}$ is referred to as a *simple* spinor field if it takes the form

$$\phi^{A_1A_2\dots A_k} = \mathrm{e}^{\psi}\xi^{A_1}\xi^{A_2}\dots\xi^{A_k}\,.$$

for some pure spinor field ξ^A, and holomorphic function ψ. The relation between simple solutions of the zrm-field equation and the existence of foliating spinors is known as the Robinson theorem [Rob61] in four dimensions, and was later generalised to even dimensions in [HM88]. Here, we give odd-dimensional versions of the theorem.

Theorem 5.17 (Robinson theorem for zrm fields) *Let ξ^A be a holomorphic pure spinor field, ψ a holomorphic function. Suppose that $\phi^{A_1A_2\dots A_k} = \mathrm{e}^{\psi}\xi^{A_1}\xi^{A_2}\dots\xi^{A_k}$ satisfies* (5.27). *Then locally*

$$\left(\xi^{a[A}\nabla_a\xi^{bB]}\right)\xi_b{}^{[C}\xi^{D]} - \frac{k-2}{2k}\left(\xi^{[A}\left(\xi^{aB]}\nabla_a\xi^{bC}\right)\xi_b{}^{D} + \xi^{[A}\left(\xi^{aB]}\nabla_a\xi^{C}\right)\xi^{D}\right) = 0\,. \tag{5.32}$$

In particular, when $k = 2$, ξ^A satisfies equation (5.17), *i.e. $\mathcal{N}^{\perp}$ is integrable.*

Suppose that ξ^A satisfies (5.17). *Then locally there exists a holomorphic function ψ such that the spinor field $\phi^{AB} = \mathrm{e}^{\psi}\xi^A\xi^B$ satisfies* (5.27). *There is the freedom of adding a holomorphic function constant along $\mathcal{N}^{\perp}$ to ψ.*

Proof. For any $\phi^{A_1 A_2 \dots A_k} = \mathrm{e}^{\psi} \xi^{A_1} \xi^{A_2} \dots \xi^{A_k}$, we have

$$\gamma^{a}{}_{B}{}^{A_1} \nabla_a \phi^{A_2 \dots A_k B} = \mathrm{e}^{\psi} \left(\xi^{A_2} \dots \xi^{A_k} \xi^{a A_1} \nabla_a \psi + (k-1) \left(\xi^{a A_1} \nabla_a \xi^{(A_2} \right) \xi^{A_3} \dots \xi^{A_k)} + \left(\nabla_a \xi^{a A_1} \right) \xi^{A_2} \dots \xi^{A_k} \right) . \tag{5.33}$$

If $\phi^{A_1 \dots A_k}$ satisfies (5.27), then we have

$$0 = \xi^{(A_2} \xi^{A_3} \dots \xi^{A_k} \xi^{a A_1)} \nabla_a \psi + (k-1) \left(\xi^{a(A_1} \nabla_a \xi^{A_2} \right) \xi^{A_3} \dots \xi^{A_k)} + \left(\nabla_a \xi^{a(A_1} \right) \xi^{A_2} \xi^{A_3} \dots \xi^{A_k)} . \tag{5.34}$$

Tensoring with $\xi^B \xi^C$ and skewing over $A_1 B$ and $A_2 C$ leads to

$$\xi^{[A} \left(\xi^{aB]} \nabla_a \xi^{[C} \right) \xi^{D]} + \xi^{[C} \left(\xi^{aD]} \nabla_a \xi^{[A} \right) \xi^{B]} = 0 , \qquad \text{i.e.} \qquad \Gamma^{(A:B)} = 0 .$$

To show that equation (5.32) also holds, we expand

$$\begin{aligned} (k-1) \left(\xi^{a(A_1} \nabla_a \xi^{A_2} \right) \xi^{A_3} \dots \xi^{A_k)} + \left(\nabla_a \xi^{a(A_1} \right) \xi^{A_2} \xi^{A_3} \dots \xi^{A_k)} & \\ = \left(- \frac{1}{4} \left(k \Gamma^{BC} - 2 \Gamma^{B:C} \right) \eta_{aB} \gamma^{a}{}_{C}{}^{(A_1} + \psi^{(A_1} \right) \xi^{A_2} \xi^{A_3} \dots \xi^{A_k)} , \end{aligned}$$

where $\psi^A \in \mathfrak{S}_{\frac{m-2}{2}}$. Since the first term on the RHS lies in $\mathfrak{S}_{\frac{m-4}{2}}$, we must have $k \Gamma^{BC} = 2 \Gamma^{B:C}$, i.e. (5.32) holds. When $k = 2$, (5.32) reduces to (5.17).

The case $k = 2$, we follow the geometrical proof given in [Eas95]. Suppose that $\mathcal{N}^{\perp}$ is integrable. Then, locally, $\mathcal{M}$ is fibered over the leaf space $\mathcal{L}$ of $\mathcal{N}^{\perp}$. Choose a holomorphic section ϕ of the tautological line bundle $\wedge^m \mathrm{T}^* \mathcal{L}$ of $\mathcal{L}$. Then, ϕ is clearly closed. Its pull-back to $\mathcal{M}$ must be orthogonal to each leaf of the foliation, i.e. it must be of the form $\phi^{AB} = \mathrm{e}^{\psi} \xi^A \xi^B$ for some holomorphic function ψ. Further ϕ is also closed, i.e. ϕ satisfy (5.27). ☐

Theorem 5.18 (Robinson theorem for co-zrm fields) *Let ξ^A be a holomorphic pure spinor field, ψ a holomorphic function. Suppose that $\phi^{A_1 A_2 \dots A_k} = \mathrm{e}^{\psi} \xi^{A_1} \xi^{A_2} \dots \xi^{A_k}$ satisfies* (5.28)*. Then locally ξ^A satisfies* (5.8)*, i.e. $\mathcal{N}$ is integrable; further, when $k > 2$, one has in addition*

$$\xi^{[A} \left(\xi^{aB]} \nabla_a \xi^{[C} \right) \xi^{D]} = 0 . \tag{5.35}$$

Suppose that ξ^A satisfies (5.8)*. Then there exists a holomorphic function ψ such that the pure spinor field $\phi^{AB} = \mathrm{e}^{\psi} \xi^A \xi^B$ satisfies* (5.28)*. Further, if ξ^A satisfies* (5.35)*, and*

$$\xi^{aA} \xi^{aB} C_{abcd} \xi^{cdC} = 0 , \tag{5.36}$$

then, for every $k > 2$, there exists a holomorphic function ψ such that the spinor field $\phi^{A_1 A_2 \dots A_k} = \mathrm{e}^{\psi} \xi^{A_1} \xi^{A_2} \dots \xi^{A_k}$ satisfies (5.28)*. In both cases, there is the freedom of adding a holomorphic function constant along $\mathcal{N}$ to ψ.*

Proof. The case $k = 2$ is similar to the proof of Theorem 5.17.

Now, for $k \geq 2$, if $\phi^{A_1 \dots A_k}$ satisfies (5.28), then equation (5.33) becomes

$$\begin{aligned} 0 = \xi^{A_3} \dots \xi^{A_k} \xi^{[A_2} \xi^{a A_1]} \nabla_a \psi + \left(\xi^{a[A_1} \nabla_a \xi^{A_2]} \right) \xi^{A_3} \dots \xi^{A_k} & \\ + (k-2) \left(\xi^{[A_2} \xi^{a A_1]} \nabla_a \xi^{(A_3} \right) \xi^{A_4} \dots \xi^{A_k)} + \left(\nabla_a \xi^{a[A_1} \right) \xi^{A_2]} \xi^{A_3} \dots \xi^{A_k} . \end{aligned} \tag{5.37}$$

Then, tensoring with ξ^B and skewing over $A_1 A_2 B$ yield (5.8). When $k > 2$, one can also tensor with ξ^B and skew over $A_3 B$, and conclude (5.35).

For the 'converse' when $k > 2$, condition (5.35) implies that

$$\begin{aligned} \xi^{aA} \nabla_a \xi^B &= \xi^A A^B + \xi^A B^B + C \xi^A \xi^B + D^A \xi^B \,, \\ \nabla_a \xi^{aA} &= E \xi^A + F^A - B^A + A^B + C \xi^A + D^A \,, \end{aligned} \tag{5.38}$$

for some functions C, E, $\mathfrak{S}^{\frac{m-2}{2}}$-valued functions B^A, D^A, F^A, and $\mathfrak{S}^{\frac{m-4}{2}}$-valued function A^A. We want to show that there exists a holomorphic function ψ such that (5.37) holds, i.e.

$$\begin{aligned} \xi^{[A} \xi^{aB]} \nabla_a \psi &= \xi^{a[A} \nabla_a \xi^{B]} + \left(\nabla_a \xi^{a[A} \right) \xi^{B]} - 2(k-2) \, \xi^{[A} D^{B]} \\ &= \xi^{[A} \left(2 \, B^{B]} - k D^{B]} - F^{B]} \right) =: \xi^{[A} \psi^{B]} \,. \end{aligned} \tag{5.39}$$

Differentiating the above equation along $\mathcal{N}$ yields the integrability condition

$$\xi^{[A} D^B \psi^{C]} = \xi^{[A} \xi^{aB} \nabla_a \psi^{C]} \,. \tag{5.40}$$

We expand the RHS of (5.40) using the expression (5.39) for ψ^A:

$$\begin{aligned} \xi^{[A} \xi^{aB} \nabla_a \psi^{C]} = (k-3) \xi^{a[A} \nabla_a \left(\xi^B D^{C]} \right) - \left(\nabla_a \xi^{a[A} \right) \xi^B D^{C]} \\ + \left(\xi^{a[A} \nabla_a \xi^B \right) \left(\nabla_b \xi^{bC]} \right) + \left(\xi^{a[A} \nabla_a \xi^B \right) \psi^{C]} + \frac{1}{8} \xi^{a[A} \xi^{bB} C_{abcd} \xi^{cdC]} \,. \end{aligned}$$

Using the identity

$$\xi^{a[A} \nabla_a \left(\xi^B D^{C]} \right) \xi^D = - \xi^{[A} D^B \left(A^{C]} + B^{C]} \right) \xi^D + \frac{1}{8} \xi^{[A} \xi^{aB} \xi^{bC]} C_{abcd} \xi^{cdD}$$

together with (5.38), (5.39) and (5.36) shows that (5.40) is indeed satisfied. ☐

Finally, we omit the proof of the following theorem, which follows roughly the one given in [HM88].

Theorem 5.19 (Non-conformally invariant Robinson theorem) *Let ξ^A be a holomorphic pure spinor field, ψ a holomorphic function. Suppose that $\phi^{A_1 A_2 \ldots A_k} = \mathrm{e}^{\psi} \xi^{A_1} \xi^{A_2} \ldots \xi^{A_k}$ satisfies* (5.29)*. Then ξ^A satisfies equation* (5.12)*, i.e. $[\Gamma(\mathcal{N}^\perp), \Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N})$.*

Suppose that ξ^A satisfies (5.12)*. There exists a holomorphic function ψ such that $\phi^{AB} = \mathrm{e}^{\psi} \xi^A \xi^B$ satisfies* (5.29)*. Suppose further that*

$$\xi^{aA} \xi^{aB} C_{abcd} \xi^{cdC} = 0 \,, \qquad \qquad \xi^{aA} \xi^{b[B} \Phi_{ab} \xi^{C]} = 0 \,. \tag{5.41}$$

Then, for every $k > 2$, there exists a hoolmorphic function ψ such that $\phi^{A_1 A_2 \ldots A_k} = \mathrm{e}^{\psi} \xi^{A_1} \xi^{A_2} \ldots \xi^{A_k}$ satisfies (5.29)*. In both cases, there is the freedom of adding a holomorphic function constant along both $\mathcal{N}$ and $\mathcal{N}^\perp$ to ψ.*

5.2.3 Pure conformal Killing spinor

Complementary to (5.26), one defines the *twistor equation*

$$\nabla_a \xi^A + \frac{1}{n} \gamma_{aB}{}^A \zeta^B = 0 \,, \tag{5.42}$$

for any holomorphic spinor field ξ^A. Here, (5.42) determines $\zeta^B = \gamma^a{}_A{}^B \nabla_a \xi^A$. A solution ξ^A will be refered to as *conformal Killing spinor*. The spinor field ζ^A can be shown to satisfy

$$\nabla_a \zeta^B + \frac{n}{2} \mathrm{P}_{ab} \gamma^b{}_A{}^B \xi^A = 0 \,, \tag{5.43}$$

where P_{ab} is the *Rho* or *Schouten* tensor, which we have defined in appendix C. These equations are conformally invariant provided that ξ^A and ζ^A transform as

$$\xi^A \mapsto \hat{\xi}^A = \xi^A \,, \qquad\qquad \zeta^A \mapsto \hat{\zeta}^A = \Omega^{-1} \left(\zeta^A + \frac{n}{2} \Upsilon_a \xi^{aA} \right) .$$

The equivalence class of spinors transforming according to this rule can be thought of as a section (ξ^A, ζ^A) of the *local twistor bundle* [PR86, BEG94] or *spin tractor bundle* [HS11], and we shall refer to such a section as a *twistor-spinor.* These are spinors for the group $\mathrm{Spin}(2m+3, \mathbb{C})$. Tracing (5.43) yields

$$\nabla_a \zeta^{aB} = - \frac{n}{4(n-1)} R \xi^B \,. \tag{5.44}$$

The following proposition is well-known, see e.g. [BJ10].

Proposition 5.20 *Let ξ^A be a conformal Killing spinor. Then*

$$C_{abcd} \xi^{cdE} = 0 \,, \qquad\qquad C_{abcd} \zeta^{bcC} - 2n A_{cab} \xi^{cE} = 0 \,, \tag{5.45}$$

where A_{abc} is the Cotton-York tensor *as defined in appendix C.*

We shall now assume that our conformal Killing spinor ξ^A is pure. Then, generically, the spinor ζ^A lies in the $\mathfrak{p}$-module $\mathfrak{S}^{\frac{m-4}{2}}$. The geometric properties of the associated γ-plane distribution essentially depends on which submodule of $\mathfrak{S}^{\frac{m-4}{2}}$ the spinor ζ^A belongs to.

Proposition 5.21 *Let ξ^A be a pure conformal Killing spinor with $\zeta^B := \gamma^a{}_A{}^B \nabla_a \xi^A$. Then ξ^A satisfies*

$$\xi^{[A} \left(\xi^{aB]} \nabla_a \xi_b{}^E \right) \xi^{b[C} \xi^{D]} + \xi^{[C} \left(\xi^{aD]} \nabla_a \xi_b{}^E \right) \xi^{b[A} \xi^{B]} = 0 \,. \tag{5.9}$$

Further, ξ^A is foliating if and only if ζ^A is pure and the respective γ-planes distribution associated to ξ^A and ζ^A intersect in a totally null $(m-1)$-plane distribution, i.e.

$$\zeta^{aA} \xi_a{}^B = - \zeta^{(A} \xi^{B)} + 3 \zeta^{[A} \xi^{B]} = \zeta^A \xi^B - 2 \xi^A \zeta^B \,. \tag{5.46}$$

Suppose ξ^A that is foliating so that ζ^A is pure and satisfies (5.46). *Then*

$$\left(\zeta^{a[A} \nabla_a \zeta^{bB} \right) \zeta_b{}^C \zeta^{D]} = 0 \,.$$

Proof. To prove that ξ^A satisfies (5.9), it suffices to contract equation (5.42) by ξ^{aA} and $\gamma^b{}_D{}^C \xi_b{}^A$. We find

$$\left(\xi^{aA} \nabla_a \xi^{bB} \right) \xi_b{}^C + \frac{1}{n} \left(\xi^{aA} \zeta_{ab}{}^B \xi^{bC} + \zeta^B \xi^A \xi^C \right) = 0 \,.$$

The second term is skew-symmetric in AC. Therefore, symmetrising over AC yields (5.9).

We now want to derive conditions on ξ^A and ζ^A for ξ^A to be foliating. In fact, it is enough to consider

$$\left(\xi^{a[A} \nabla_a \xi^B \right) \xi^{C]} = 0 \,. \tag{5.8}$$

Contracting (5.42) with ξ^{aA} yields immediately the condition $\xi^{a[A} \zeta_a{}^B \xi^{C]} = 0$, which is equivalent to (5.46) – the factors can be obtained by means of Proposition 2.9. Assuming now that ξ^A and ζ^A satisfy (5.46), we compute

$$\xi^{aB} \zeta_a{}^A \gamma^b{}_A{}^C \xi_b{}^E = \left(\zeta^A \xi^B - 2 \xi^A \zeta^B \right) \gamma^b{}_A{}^C \xi_b{}^E = \zeta^C \xi^E \xi^B - 2 \xi^C \zeta^E \xi^B + 2 \xi^C \xi^E \zeta^B \,,$$

from which we deduce the remaining condition (5.7).

Finally, if ξ^A and ζ^A are pure and satisfy (5.46), contracting equation (5.43) with ζ^{aA} and $\gamma^b{}_D{}^C \zeta_b{}^A$ leads to

$$\left(\zeta^{aA} \nabla_a \zeta^{bB} \right) \zeta_b{}^C - \frac{1}{2} \zeta^{aA} \mathrm{P}_{ab} \xi^{bB} \zeta^C + 2 \zeta^{aA} \mathrm{P}_{ab} \zeta^{b[B} \xi^{C]} = 0 \,,$$

and the result follows by symmetry considerations. □

Remark 5.22 There are two ways of looking at Proposition 5.21. One is to say that the conformal Killing spinor ξ^A is foliating if and only if ζ^A lies in the $\mathfrak{p}$-module $\mathfrak{S}^{\frac{m-2}{2}}$. The other is to think of the twistor-spinor (ξ^A, ζ^A) as *pure* as a spinor for the conformal group.

5.2.4 Pure Killing spinors

A holomorphic spinor field ξ^A which is both a solution to the twistor equation (5.42) and an eigenspinor of the Dirac operator, i.e.

$$\gamma^a{}_B{}^C \nabla_a \xi^B = \lambda \xi^C \,,$$

for some smooth function λ on $\mathcal{M}$ is known as a *Killing spinor.* Otherwise put, ξ^A satisfies the Killing equation

$$\nabla_a \xi^A + \lambda \frac{1}{n} \xi_a{}^A = 0 \,. \tag{5.47}$$

In the case where ξ^A is pure, then ξ^A is a Killing spinor if and only if $\zeta^A \in \mathfrak{S}^{\frac{m}{2}}$. This can be seen as an extension of Proposition 5.21: at any point of $\mathcal{M}$, ζ^A takes values in $\mathfrak{S}^{\frac{m-4}{2}}$. Then ξ^A is foliating if and only if locally $\zeta^A \in \mathfrak{S}^{\frac{m-2}{2}}$.

The Killing spinor equation (5.47) is clearly not conformally invariant, and one will expect non-conformally invariant integrability conditions and other geometric properties.

Proposition 5.23 *Let ξ^A be a pure Killing spinor. Then*

$$C_{abcd} \xi^{cdD} = 0 \,, \qquad \xi^{aA} A_{abc} = 0$$
$${}^{\mathfrak{F}}\Pi^1_0(\Phi) = 0 \,.$$

Further, its eigenvalue λ satisfies

$$\xi^{aA} \nabla_a \lambda = - \left(\lambda^2 + \frac{n}{4(n-1)} R \right) \xi^A \,,$$

i.e. λ is constant along the γ-plane distribution defined by ξ^A.

Proof. This is a special case of (5.42), (5.43), (5.44), and (5.45) with $\zeta^A = \lambda \xi^A$. □

This equation is also valid in even dimensions, but must necessarily involve non-irreducible Dirac spinors. Since there is no chirality in odd dimensions, one is free to restrict the Killing equation to pure spinor fields. The relation to the foliation condition is given by the following lemma, whose proof is left for the reader.

Proposition 5.24 *Let ξ^A be a pure conformal Killing spinor. Then ξ^A is a Killing spinor if and only if*

$$\left(\nabla_a \xi^{b[A} \right) \xi_b{}^B \xi^{C]} = 0 \,, \qquad \left(\xi^{aC} \nabla_a \xi^{[A} \right) \xi^{B]} = 0 \,.$$

In particular, ξ^A is strongly foliating, i.e. $[\Gamma(\mathcal{N}^\perp), \Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N})$.

The question of when a pure conformal Killing spinor is locally a parallel spinor was dealt with in even dimensions in [TC12b]. In odd dimensions, things are very similar, and the next result will be stated without proof.

Proposition 5.25 *Let ξ^A be a foliating pure conformal Killing spinor, and suppose that the metric can be rescaled so that the Schouten tensor satisfies $\mathrm{P}_{ab} \xi^{aA} = 0$. Then there exists a conformal rescaling such that ξ^A is locally parallel.*

Remark 5.26 Non-foliating twistor-spinors are pertinent to the theory of differential equations. Using the method of equivalence, Cartan [Car10] showed how to encode the invariance properties of certain ODEs of Monge type in terms of a $(2,3,5)$-distribution, i.e. a rank-2 distribution $\mathcal{N}$ on a 5-dimensional smooth manifold, generating the tangent bundle, i.e. $[\mathcal{N},\mathcal{N}]$ is a rank-3 distribution, and $[\mathcal{N},[\mathcal{N},\mathcal{N}]]$ is the full tangent bundle. This is more invariant expressed as a G_2-principle bundle equipped with a Cartan connection. To this $(2,3,5)$-distribution, Nurowski [Nur05] was then able to associate a split-signature five-dimensional conformal structure, with respect to which $\mathcal{N}$ is totally null, with orthogonal complement $[\mathcal{N},\mathcal{N}]$. The general theory, expounded in the language of parabolic geometries, is given in [ČS09a, HS09, HS11]. In particular, the conformal holomony of such manifolds is characterised by being contained in G_2, or equivalently, by the existence of a real twistor-spinor, *generic* in the sense that $\xi^A \zeta_A \neq 0$.

Flat case For a $(2m+1)$-dimensional conformally flat manifold $\mathbb{M}$, the twistor equation can be integrated explicitly

$$\xi^A = \mathring{\xi}^A + x^a \gamma_{aB}{}^A \mathring{\zeta}^B \,, \qquad\qquad \zeta^B = \mathring{\zeta}^B \,,$$

for some constant spinors $\mathring{\xi}^A$, $\mathring{\zeta}^B$. The space of solutions $(\mathring{\xi}^A, \mathring{\zeta}^B)$ is thus a 2^m-dimensional vector space. The solutions of $(\mathring{\xi}^A, \mathring{\zeta}^B)$ distinguished by $\mathring{\xi}^A$, $\mathring{\zeta}^B$ pure, and satisfying the constraint (5.46), has then a nice geometrical interpretation. Its projectivation is the space of all γ-planes in the compactification of $\mathbb{M}$, also known as the *twistor space* of $\mathbb{M}$.

Relation to the Goldberg-Sachs theorem In [TC12a], a higher-dimensional version of the Goldberg-Sachs theorem is given, based on a coarser version of the present classification of the Weyl tensor and Cotton-York tensor. In particular, suppose the Weyl tensor and the Cotton-York tensor satisfies the conditions

$$\xi^{aA} \xi^{bB} \xi^{c[C} C_{abcd} \xi^{D]} = 0 \,, \qquad\qquad \xi^{[A} \xi^{aB]} \xi^{bC} \xi^{cD} A_{abc} = 0 \,. \tag{5.48}$$

Suppose further that the Weyl tensor is otherwise generic. Then the pure spinor field $\xi^{A'}$ is foliating.

As in the even-dimensional case [TC12b], Proposition 5.21 provides an example of a non-foliating pure spinor field which satisfies (5.48), but which however violates the genericity assumption by virtue of Proposition 5.20. While the spinor is not foliating, its intrinsic torsion satisfies a weaker condition. This motivates the following conjecture improving the author's work [TC12a].

Conjecture 5.27 *Let ξ^A be a holomorphic pure spinor field on a $(2m+1)$-dimensional Einstein complex Riemannian manifold $(\mathcal{M}, g)$ such that the Weyl tensor satisfies $\xi^{aA} \xi^{bB} \xi^{c[C} C_{abcd} \xi^{D]} = 0$. Then ξ^A satisfies*

$$\xi^{[A} \left(\xi^{aB]} \nabla_a \xi_b{}^E \right) \xi^{b[C} \xi^{D]} + \xi^{[C} \left(\xi^{aD]} \nabla_a \xi_b{}^E \right) \xi^{b[A} \xi^{B]} = 0 \,. \tag{5.9}$$

Weaker conditions such as (5.6) may well be possible too, but an investigation of the veracity of the above conjecture is beyond the scope of this article.

A Spinorial descriptions of curvature tensors

A.1 Projection maps

In this appendix, we give explicit formulae describing the various irreducibles $\mathfrak{p}$-modules of the spaces $\mathfrak{F}$, $\mathfrak{A}$ and $\mathfrak{C}$ as kernels of maps ${}^{\mathfrak{F}}\Pi_i^j$, ${}^{\mathfrak{A}}\Pi_i^j$ and ${}^{\mathfrak{C}}\Pi_i^j$, which on restriction to the $\mathfrak{p}$-modules $\mathrm{gr}_i(\mathfrak{F})$, $\mathrm{gr}_i(\mathfrak{A})$ and $\mathrm{gr}_i(\mathfrak{C})$ project into the ireeducible $\mathfrak{p}$-modules $\mathfrak{F}_i^j$, $\mathfrak{A}_i^j$ and $\mathfrak{C}_i^j$.

Tracefree Ricci tensor For $\Phi_{ab} \in \mathfrak{F}$, define

$$
\begin{aligned}
{}^{\mathfrak{F}}\Pi^0_{-2}(\Phi) &:= \xi^{[A}\,\xi^{a\,B]}\Phi_{ab}\xi^{b[C}\,\xi^{\,D]}\,, &\qquad {}^{\mathfrak{F}}\Pi^0_{-1}(\Phi) &:= \xi^{[A}\,\xi^{a\,B]}\Phi_{ab}\xi^{bC}\,,\\
{}^{\mathfrak{F}}\Pi^0_{0}(\Phi) &:= \xi^{aA}\xi^{bB}\Phi_{ab}\,, & {}^{\mathfrak{F}}\Pi^1_{0}(\Phi) &:= \xi^{[A}\,\xi^{a\,B]}\Phi_{ab} + \frac{1}{n-1}\gamma_{bC}{}^{[A}\,\xi^{c\,B]}\xi^{dC}\Phi_{cd}\,,\\
{}^{\mathfrak{F}}\Pi^0_{1}(\Phi) &:= \xi^{aA}\Phi_{ab}\,,
\end{aligned}
$$

Cotton-York tensor For $A_{abc} \in \mathfrak{A}$, define

$$
\begin{aligned}
&\Pi^0_{-3}(A) := \xi^{[A}\,\xi^{a\,B]}\xi^{b[C}\,\xi^{cD}\xi^{\,E]}A_{abc}\,,\\
&\Pi^0_{-2}(A) := \xi^{a(A}\xi^{bB)}\xi^{cC}A_{abc} \quad (\mathrm{mod}\ \xi^{(A}\alpha^{B)C})\,, \qquad \Pi^1_{-2}(A) := \xi^{a[A}\xi^{bB]}\xi^{cC}A_{abc} \quad (\mathrm{mod}\ \xi^{[A}\alpha^{B]C})\,,
\end{aligned}
$$

$$
\begin{aligned}
&\Pi^3_{-1}(A) := \xi^{C}\xi^{a(A}\xi^{bB)}A_{abc} + \frac{3}{2(n+1)}\xi^{C}\xi^{a(A|}\xi^{bdD}A_{abd}\gamma_{cD}{}^{|B)} - \frac{1}{n+1}\xi^{aD}\xi^{bC}\xi^{d(A|}A_{abd}\gamma_{cD}{}^{|B)} \quad (\mathrm{mod}\ \xi^{C}\xi^{(A}\phi^{B)}{}_{c})\,,\\
&\Pi^2_{-1}(A) := \xi^{C}\xi^{a[A}\xi^{bB]}A_{abc} + \frac{1}{2(n-3)}\xi^{C}\xi^{a[A|}\xi^{bdD}A_{abd}\gamma_{cD}{}^{|B]} - \frac{1}{n-3}\xi^{aD}\xi^{bC}\xi^{d[A|}A_{abd}\gamma_{cD}{}^{|B]} \quad (\mathrm{mod}\ \xi^{C}\xi^{[A}\phi^{B]}{}_{c})\,,\\
&\Pi^1_{-1}(A) := \xi^{aA}\xi^{bB}\xi^{cC}A_{abc}\,,\\
&\Pi^0_{-1}(A) := \xi^{[A}\,\xi^{a\,B]}\xi^{bcC}A_{abc} - \frac{1}{n-2}\xi^{aC}\xi^{bA}\xi^{cB}A_{abc}\,,
\end{aligned}
$$

$$
\begin{aligned}
&\Pi^2_{0}(A) := \xi^{a(A}\xi^{bB)}A_{abc} \quad (\mathrm{mod}\ \xi^{(A}\phi^{B)}{}_{c})\,, \qquad \Pi^1_{0}(A) := \xi^{a[A}\xi^{bB]}A_{abc} \quad (\mathrm{mod}\ \xi^{[A}\phi^{B]}{}_{c})\,,\\
&\Pi^0_{0}(A) := \xi^{aA}\xi^{bcB}A_{abc}\,,
\end{aligned}
$$

$$
\begin{aligned}
&\Pi^3_{1}(A) := \xi^{cC}A_{(ab)c}\xi^{D} - \frac{3}{2(n+1)}\gamma_{(aB}{}^{C}A_{b)cd}\xi^{cdB}\xi^{D} + \frac{1}{n+1}\xi^{cC}\xi^{dB}A_{cd(a}\gamma_{b)B}{}^{D}\,,\\
&\Pi^2_{1}(A) := \xi^{cC}A_{[ab]c}\xi^{D} - \frac{1}{2(n-3)}\gamma_{(aB}{}^{C}A_{b)cd}\xi^{cdB}\xi^{D} + \frac{1}{n-3}\xi^{cC}\xi^{dB}A_{cd[a}\gamma_{b]B}{}^{D}\,,
\end{aligned}
$$

$$
\Pi^1_{1}(A) := \xi^{aA}\xi^{bB}A_{abc}\,, \qquad \Pi^0_{1}(A) := \xi^{bcC}A_{abc}\xi^{D} + \frac{2}{n-2}\xi^{bB}\xi^{bC}A_{bca}\,,
$$

$$
\Pi^0_{2}(A) := \xi^{cC}A_{[ab]c}\,, \qquad \Pi^1_{2}(A) := \xi^{cC}A_{(ab)c}\,.
$$

Weyl tensor For $C_{abcd} \in \mathfrak{C}$, define

$$
\begin{aligned}
&{}^{\mathfrak{C}}\Pi^0_{-4}(C) := \xi^{[A}\,\xi^{aB}\xi^{b\,C]}C_{abcd}\xi^{c[D}\,\xi^{dE}\xi^{\,F]}\,,\\
&{}^{\mathfrak{C}}\Pi^0_{-3}(C) := \xi^{[A}\,\xi^{aB}\xi^{b\,C]}C_{abcd}\xi^{cD}\xi^{dE}\,,
\end{aligned}
$$

$$
{}^{\mathfrak{C}}\Pi^0_{-2}(C) := \xi^{[A}\,\xi^{aB}\xi^{b\,C]}C_{abcd}\xi^{cdD}\,, \qquad {}^{\mathfrak{C}}\Pi^1_{-2}(C) := \xi^{aA}\xi^{bB}C_{abcd}\xi^{cC}\xi^{dD}\,,
$$

$$
\begin{aligned}
{}^{\mathfrak{C}}\Pi^2_{-2}(C) := \xi^{aA}\xi^{bB}C_{abcd}\xi^{cC}\xi^{D} &+ \frac{1}{n+1}\left(\xi^{aA}\xi^{bB}C_{abce}\xi^{ceE}\gamma_{dE}{}^{C} - \xi^{aC}\xi^{b[A|}C_{abce}\xi^{ceE}\gamma_{dE}{}^{|B]}\right)\xi^{D}\\
&- \frac{1}{n-3}\gamma_{dE}{}^{[A|}\xi^{aE}\xi^{b|B]}C_{abde}\xi^{dC}\xi^{eD} \quad (\mathrm{mod}\ \xi^{[A}\alpha_{d}{}^{B]C}\xi^{D} - \xi^{C}\alpha_{d}{}^{AB}\xi^{D})\,,
\end{aligned}
$$

$$
\begin{aligned}
{}^{\mathfrak{C}}\Pi^0_{-1}(C) &:= \xi^{aA}\xi^{bB}C_{abcd}\xi^{cdC} \,, \\
{}^{\mathfrak{C}}\Pi^1_{-1}(C) &:= \xi^{aA}\xi^{b[B}C_{abcd}\xi^{cC]} + \frac{1}{n-3}\gamma_{dE}{}^{[B|}\xi^{aE}\xi^{b|C]}\xi^{ceA}C_{abce} \quad (\mathrm{mod}\ \xi^A\xi^{[B}\alpha_d{}^{C]}) \,, \\
{}^{\mathfrak{C}}\Pi^2_{-1}(C) &:= \xi^{aA}\xi^{b(B}C_{abcd}\xi^{cC)} + \frac{3}{n+1}\gamma_{dE}{}^{(B|}\xi^{aE}\xi^{b|C)}\xi^{ceA}C_{abce} \quad (\mathrm{mod}\ \xi^A\xi^{(B}\alpha_d{}^{C)}) \,,
\end{aligned}
$$

$$
{}^{\mathfrak{C}}\Pi^0_0(C) := \xi^{abA}C_{abcd}\xi^{cdB} \,, \qquad {}^{\mathfrak{C}}\Pi^1_0(C) := \xi^{abA}C_{abcd}\xi^{d[C}\xi^{D]} - \frac{1}{n-1}\xi^{abA}\xi^{deE}C_{abde}\gamma_{cE}{}^{[C}\xi^{D]} \,,
$$

$$
{}^{\mathfrak{C}}\Pi^2_0(C) := \xi^{aA}\xi^{bB}C_{abcd}\xi^{cC} \,,
$$

$$
\begin{aligned}
{}^{\mathfrak{C}}\Pi^3_0(C) := \xi^{aA}C_{a[bc]d}\xi^{dD}\xi^C &+ \frac{1}{n-5}\left(\gamma_{[b|E}{}^{[A|}\xi^{aeE}C_{ae|c]d}\xi^{d|D]}\xi^C - \xi^{aC}\xi^{e[A}C_{ae[b|f}\xi^{fE}\gamma_{c]E}{}^{D]}\right) \\
&- \frac{1}{2(n-3)(n-5)}\gamma_{[b|E}{}^{[A}\xi^{aeE}C_{aedf}\xi^{dfF}\gamma_{|c]F}{}^{D]}\xi^C \quad (\mathrm{mod}\ \xi^{[A}\alpha_{bc}{}^{D]}\xi^C) \,,
\end{aligned}
$$

$$
\begin{aligned}
{}^{\mathfrak{C}}\Pi^4_0(C) := \xi^{aA}C_{a(bc)d}\xi^{dD}\xi^C &+ \frac{3}{n+3}\left(\gamma_{(b|E}{}^{(A|}\xi^{aeE}C_{ae|c)d}\xi^{d|D)}\xi^C - \xi^{aC}\xi^{e(A|}C_{ae(b|f}\xi^{fE}\gamma_{c)E}{}^{|D)}\right) \\
&- \frac{3}{2(n+1)(n+3)}\gamma_{(b|E}{}^{(A}\xi^{aeE}C_{aedf}\xi^{dfF}\gamma_{|c)F}{}^{D)}\xi^C \quad (\mathrm{mod}\ \xi^{(A}\alpha_{bc}{}^{D)}\xi^C) \,,
\end{aligned}
$$

$$
{}^{\mathfrak{C}}\Pi^0_1(C) := \xi^{abB}C_{abcd}\xi^{dC} \,, \qquad {}^{\mathfrak{C}}\Pi^1_1(C) := \xi^{aA}C_{a[bc]d}\xi^{dD} + \frac{1}{n-3}\gamma_{[b|E}{}^{[A|}\xi^{aeE}C_{ae|c]d}\xi^{d|D]} \,,
$$

$$
{}^{\mathfrak{C}}\Pi^2_1(C) := \xi^{aA}C_{a(bc)d}\xi^{dD} + \frac{3}{n+1}\gamma_{(b|E}{}^{(A|}\xi^{aeE}C_{ae|c)d}\xi^{d|D)} \quad (\mathrm{mod}\ \xi^A\xi^D\alpha_{bc}) \,,
$$

$$
{}^{\mathfrak{C}}\Pi^0_2(C) := \xi^{abA}C_{abcd} \,, \qquad {}^{\mathfrak{C}}\Pi^1_2(C) := \xi^{aA}C_{a(bc)d}\xi^{dB} \,,
$$

$$
\begin{aligned}
{}^{\mathfrak{C}}\Pi^2_2(C) := \xi^D\xi^{aA}C_{abcd} &+ \frac{1}{n+1}\left(\gamma_{bE}{}^A\xi^{aeE}C_{aecd} - \gamma_{[cE}{}^A\xi^{aeE}C_{ae|d]b}\right)\xi^D \\
&+ \frac{1}{2(n-3)}\left(\gamma_{[c|E}{}^A\xi^{aE}C_{a|d]be}\xi^{eD}\right) \quad (\mathrm{mod}\ \xi^A\xi^D\alpha_{bc}) \,,
\end{aligned}
$$

$$
{}^{\mathfrak{C}}\Pi^0_3(C) := \xi^{aA}C_{abcd} \,,
$$

A.2 Spinorial representatives

It is often more convenient to work with a particular choice of grading adapted to the filtration (2.4). In particular, the irreducibles $\mathfrak{p}$-modules $\mathfrak{F}_i^j$, $\mathfrak{A}_i^j$ and $\mathfrak{C}_i^j$ are isomorphic to irreducibles $\mathfrak{g}_0$-modules $\breve{\mathfrak{F}}_i^j$, $\breve{\mathfrak{A}}_i^j$ and $\breve{\mathfrak{C}}_i^j$ respectively, and one can give explicit spinorial 'representatives' for elements of those modules.

In what follows, ξ^A and η_A are spinors annihiling $\mathfrak{V}_1$ and $\mathfrak{V}_{-1}$ and satisfying $\xi^A\eta_A = -\frac{1}{2}$, and

$$
u_a = 2\,\xi^{aA}\eta_A \,, \qquad h_{ab} = g_{ab} + u_a u_b \,, \qquad \omega_{ab} = -2\,\xi_{[a}{}^A\eta_{b]A} \,,
$$

as already defined in (2.15), (2.16) and (2.20) respectively. Further, any upstairs and downstairs upper case Roman index refers to the $\mathfrak{g}_0$-modules $\mathfrak{S}_{\frac{m-2}{2}}$ and $\mathfrak{S}_{-\frac{m-2}{2}}$ respectively. Finally, an element

$$
\sigma_{A\ldots C}{}^{D\ldots F} \quad \in \quad \mathfrak{S}_{-\frac{m-2}{2}} \otimes \ldots \otimes \mathfrak{S}_{-\frac{m-2}{2}} \otimes \mathfrak{S}_{\frac{m-2}{2}} \otimes \ldots \otimes \mathfrak{S}_{\frac{m-2}{2}}
$$

will be referred to as *(totally) tracefree*, if the contraction of any pair of indices with the identity element I_A^B defined by (2.13) vanishes, e.g. $\sigma_A{}^B I_B^A = 0$.

Tracefree Ricci tensor Let $\Phi_{ab} \in \mathfrak{F}$. Then

- $\Phi_{ab} \in \breve{\mathfrak{F}}^1_0$ if and only if $\Phi_{ab} = \xi_{(a}{}^{A} \eta_{b)B} \Phi_{A}{}^{B}$ for some tracefree $\Phi_{A}{}^{B}$;
- $\Phi_{ab} \in \breve{\mathfrak{F}}^0_0$ if and only if $\Phi_{ab} = \Phi \left(u_a u_b + \frac{1}{n-1} h_{ab} \right)$ for some complex Φ;
- $\Phi_{ab} \in \breve{\mathfrak{F}}^0_1$ if and only if $\Phi_{ab} = \xi_{[a}{}^{A} \Phi_A u_{b]}$ for some Φ_A;
- $\Phi_{ab} \in \breve{\mathfrak{F}}^0_2$ if and only if $\Phi_{ab} = \xi_a{}^A \xi_b{}^B \Phi_{AB}$ for some $\Phi_{AB} = \Phi_{(AB)}$, and similarly for an element of $\breve{\mathfrak{F}}^0_{-2}$ by substituting ξ^A for η_A and changing the index structure in the obvious way.

Cotton-York tensor Let $A_{abc} \in \mathfrak{A}$. Then

- $A_{abc} \in \breve{\mathfrak{A}}^0_0$ if and only if $A_{abc} = a \left(u_a \omega_{bc} - u_{[b} \omega_{c]a} \right)$ for some complex a;
- $A_{abc} \in \breve{\mathfrak{A}}^1_0$ if and only if $A_{abc} = u_a A_{bc} - u_{[b} A_{c]a}$ where $A_{ab} = \xi_{[a}{}^{A} \eta_{b]B} A_{A}{}^{B}$ for some tracefree $A_{A}{}^{B}$;
- $A_{abc} \in \breve{\mathfrak{A}}^2_0$ if and only if $A_{abc} = A_{a[b} u_{c]}$ where $A_{ab} = \xi_{(a}{}^{A} \eta_{b)B} A_{A}{}^{B}$ for some tracefree $A_{A}{}^{B}$;
- $A_{abc} \in \breve{\mathfrak{A}}^0_1$ if and only if $A_{abc} = A_a \omega_{bc} - A_{[b} \omega_{c]a} + \frac{3}{n-1} g_{a[b} \omega_{c]d} A^d$ where $A_c = \xi_c{}^C A_C$ for some A_A;
- $A_{abc} \in \breve{\mathfrak{A}}^1_1$ if and only if $A_{abc} = 2 u_a u_{[b} A_{c]} + \frac{2}{n-2} h_{a[b} A_{c]}$ where $A_a = \xi_a{}^A A_A$ for some A_A;
- $A_{abc} \in \breve{\mathfrak{A}}^2_1$ if and only if $A_{abc} = \xi_b{}^A \xi_c{}^B \eta_{aC} A_{AB}{}^{C} - \xi_a{}^A \xi_{[b}{}^{B} \eta_{c]C} A_{AB}{}^{C}$ for some tracefree $A_{AB}{}^{C} = A_{[AB]}{}^{C}$;
- $A_{abc} \in \breve{\mathfrak{A}}^3_1$ if and only if $A_{abc} = \xi_a{}^A \xi_{[b}{}^{B} \eta_{c]C} A_{AB}{}^{C}$ for some tracefree $A_{AB}{}^{C} = A_{(AB)}{}^{C}$;
- $A_{abc} \in \breve{\mathfrak{A}}^0_2$ if and only if $A_{abc} = u_a A_{bc} - u_{[b} A_{c]a}$ where $A_{ab} = \xi_a{}^A \xi_b{}^B A_{AB}$ fo some $A_{AB} = A_{[AB]}$;
- $A_{abc} \in \breve{\mathfrak{A}}^1_2$ if and only if $A_{abc} = A_{a[b} u_{c]}$ where $A_{ab} = \xi_a{}^A \xi_b{}^B A_{AB}$ for some $A_{AB} = A_{(AB)}$;
- $A_{abc} \in \breve{\mathfrak{A}}^0_3$ if and only if $A_{abc} = \xi_a{}^A \xi_b{}^B \xi_c{}^C A_{ABC}$ for some $A_{ABC} = A_{A[BC]}$ satisfying $A_{[ABC]} = 0$.

Using the duality $(\breve{\mathfrak{A}}^j_{-i})^* \cong \breve{\mathfrak{A}}^j_i$, spinorial decompositions of elements of $\breve{\mathfrak{A}}^j_{-i}$ for $i = 1, 2, 3$ can be obtained by interchanging ξ^A and η_A in the obvious way and notation.

Weyl tensor Let $C_{abcd} \in \mathfrak{C}$. Then

- $C_{abcd} \in \breve{\mathfrak{C}}^0_0$ if and only if

  $$C_{abcd} = c \left(2\omega_{ab}\omega_{cd} - 2\omega_{a[c} \omega_{d]b} + \frac{6}{n-2} h_{a[c} h_{d]b} \right) ;$$

  for some complex c;
- $C_{abcd} \in \breve{\mathfrak{C}}^1_0$ if and only if

  $$C_{abcd} = \omega_{ab} \Psi_{cd} + \Psi_{ab}\omega_{cd} - 2\omega_{[a\,|[c} \Psi_{d]|\,b]} - \frac{6}{n-3} \left(h_{[a\,|[c} \omega_{d]}{}^{e} \Psi_{|\,b]e} + h_{[c\,|[a} \omega_{b]}{}^{e} \Psi_{|\,d]e} \right) .$$

  where $\Psi_{cd} := 2\xi_{[c}{}^{C} \eta_{d]D} \Psi_{C}{}^{D}$ for some tracefree $\Psi_{C}{}^{D}$;

- $C_{abcd} \in \breve{\mathfrak{C}}^2_0$ if and only if
$$C_{abcd} = u_{[a}\,\Psi_{b][c}\,u_{d]} - \frac{1}{n-3} h_{[a\,|[c}\,\Psi_{d]|\,b]}\,,$$
where $\Psi_{cd} := 2\xi_{(c}{}^{C}\eta_{d)D}\Psi_{C}{}^{D}$ for some tracefree $\Psi_{C}{}^{D}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^3_0$ if and only if
$$C_{abcd} = \xi_a{}^A\xi_b{}^B\eta_{cC}\eta_{dD}\Psi_{AB}{}^{CD} + \xi_c{}^A\xi_d{}^B\eta_{aC}\eta_{bD}\Psi_{AB}{}^{CD} - 2\xi_{[a\,|}{}^{A}\xi_{[c}{}^{C}\eta_{d]|D}\eta_{b]B}\Psi_{AC}{}^{DB}\,,$$
for some tracefree $\Psi_{AC}{}^{DB} = \Psi_{[AC]}{}^{[DB]}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^4_0$ if and only if
$$C_{abcd} = \xi_{[a\,|}{}^{A}\xi_{[c}{}^{C}\eta_{d]|D}\eta_{b]B}\Psi_{AC}{}^{DB}\,,$$
for some tracefree $\Psi_{AC}{}^{DB} = \Psi_{(AC)}{}^{(DB)}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^0_1$ if and only if
$$\begin{aligned} C_{abcd} = \omega_{ab}\Psi_{[c}\,u_{d]} + \omega_{cd}\Psi_{[a}\,u_{b]} - \omega_{[a\,|[c}\,\Psi_{d]}u_{|\,b]} - \omega_{[c\,|[a}\,\Psi_{b]}u_{|\,d]} \\ + \frac{3}{n-2}\left(g_{[a\,|[c}\,u_{d]}\omega_{|\,b]}{}^{e}\Psi_e + g_{[c\,|[a}\,u_{b]}\omega_{|\,d]}{}^{e}\Psi_e\right)\,. \end{aligned}$$
where $\Psi_a = \xi_a{}^A\Psi_A$ for some Ψ_A;
- $C_{abcd} \in \breve{\mathfrak{C}}^1_1$ if and only if
$$C_{abcd} = u_{[a}\,\Psi_{b]cd} + u_{[c}\,\Psi_{d]ab}\,,$$
where $\Psi_{abc} = \xi_a{}^A\xi_b{}^B\eta_{cC}\Psi_{BC}{}^{D} - \xi_c{}^A\xi_{[a}{}^{B}\eta_{b]C}\Psi_{BC}{}^{D}$ for some tracefree $\Psi_{AB}{}^{D} = \Psi_{[AB]}{}^{D}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^2_1$ if and only if
$$C_{abcd} = u_{[a}\,\Psi_{b]cd} + u_{[c}\,\Psi_{d]ab}\,,$$
where $\Psi_{abc} = \xi_c{}^A\xi_{[a}{}^{B}\eta_{b]C}\Psi_{BC}{}^{D}$ for some tracefree $\Psi_{AB}{}^{D} = \Psi_{(AB)}{}^{D}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^0_2$ if and only if
$$C_{abcd} = \omega_{ab}\Psi_{cd} + \Psi_{ab}\omega_{cd} - 2\omega_{[a\,|[c}\,\Psi_{d]|\,b]} - \frac{6}{n-2}\left(g_{[a\,|[c}\,\omega_{d]}{}^{e}\Psi_{|\,b]e} + g_{[c\,|[a}\,\omega_{b]}{}^{e}\Psi_{|\,d]e}\right)\,.$$
where $\Psi_{ab} := \xi_a{}^A\xi_b{}^B\Psi_{AB}$ for some $\Psi_{CD} = \Psi_{[CD]}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^1_2$ if and only if
$$C_{abcd} = u_{[a}\,\Psi_{b][c}\,u_{d]} - \frac{1}{n-3} h_{[a\,|[c}\,\Psi_{d]|\,b]}\,,$$
where $\Psi_{cd} := \xi_c{}^C\xi_d{}^D\Psi_{CD}$ for some $\Psi_{AB} = \Psi_{(AB)}$;
- $C_{abcd} \in \breve{\mathfrak{C}}^2_2$ if and only if
$$C_{abcd} = \xi_a{}^A\xi_b{}^B\xi_{[c}{}^{C}\eta_{d]D}\Psi_{ABC}{}^{D} + \xi_c{}^A\xi_d{}^B\xi_{[a}{}^{C}\eta_{b]D}\Psi_{ABC}{}^{D}\,,$$
for some $\Psi_{ABC} = \Psi_{[AB]C}$ satisfying $\Psi_{[ABC]} = 0$;

- $C_{abcd} \in \breve{\mathfrak{C}}^0_3$ if and only if
$$C_{abcd} = u_{[a}\,\Psi_{\,b]cd} + u_{[c}\,\Psi_{\,d]ab}\,,$$
where $\Psi_{abc} = \xi_a{}^A \xi_b{}^B \xi_c{}^C \Psi_{ABC}$ for some $\Psi_{ABC} = \Psi_{[AB]C}$ satisfying $\Psi_{[ABC]} = 0$;
- $C_{abcd} \in \breve{\mathfrak{C}}^0_4$ if and only if
$$C_{abcd} = \xi_a{}^A \xi_b{}^B \xi_c{}^C \xi_d{}^D \Psi_{ABCD}\,,$$
for some $\Psi_{ABCD} = \Psi_{[AB][CD]}$ satisfying $\Psi_{[ABC]D} = 0$.

Using the duality $(\breve{\mathfrak{C}}^j_{-i})^* \cong \breve{\mathfrak{C}}^j_i$, spinorial decompositions of elements of $\breve{\mathfrak{C}}^j_{-i}$ for $i = 1, 2, 3$ can be obtained by interchanging ξ^A and η_A in the obvious way and notation.

B Spinor calculus in three and five dimensions

In this appendix, we give a brief description of spinor calculus in three and five dimensions emphasising its relation to the classification of tensors with respect to a ‘principal spinor’.

B.1 Three dimensions

Let $(\mathfrak{V}, g)$ be a three-dimensional complex vector space equipped with a non-degenerate symmetric bilinear form. The double cover of the special orthogonal groups $\mathrm{SO}(3,\mathbb{C})$ is the complex special linear group $\mathrm{SL}(2,\mathbb{C})$. Thus, the spin space $\mathfrak{S}$ and its dual space are two-dimensional complex vector spaces equipped with non-degenerate symplectic forms ε_{AB} and ε^{AB} respectively. By the two-dimensionality of $\mathfrak{S}$, it thus follows that, as in the case of the two-spinor calculus in four dimensions,[4] any irreducible spinor representations must be totally symmetric since skew-symmetry can be eliminated by means of the volume forms ε_{AB} and ε^{AB}. We take the convention that $\varepsilon_{AC}\varepsilon^{BC} = \delta_A^B$ and indices are lowered and raised as follows
$$\xi^A \varepsilon_{AB} = \xi_B\,, \qquad\qquad \varepsilon^{AB}\xi_B = \xi^A\,.$$

We can convert tensorial quantities into spinorial ones by means of the normalised Van der Waerden γ-matrices $\frac{1}{\sqrt{2}}\gamma_a{}^{AB}$, which are symmetric in their spinor indices, and satisfy the identity
$$\gamma_{aA}{}^{B}\gamma^{a}{}_{C}{}^{D} = -\delta_A^D\delta_C^B + \varepsilon_{AC}\varepsilon^{BD}\,, \qquad \text{i.e.} \qquad \gamma_{aAB}\gamma^{a}{}_{CD} = -2\varepsilon_{A(C}\,\varepsilon_{\,D)B}\,.$$

In particular, we have that $\mathfrak{V} \cong \wedge^2\mathfrak{V} \cong \odot^2\mathfrak{S}$ where we have made use of Hodge duality, so that two-forms are also represented by symmetric spinors of valence 2. Further, since it is well known that there is no Weyl tensor in dimension three, the only irreducible curvature tensors that remain are the tracefree Ricci tensor $\Phi_{ABCD} = \Phi_{(ABCD)}$ and the Cotton-York tensor $A_{ABCD} = A_{(ABCD)}$. In particular, the spaces $\mathfrak{F}$ and $\mathfrak{A}$ of tracefree Ricci tensors and of Cotton-York tensors are isomorphic to one another.

Principal spinors Fixing a spinor ξ^A induces a filtration $\mathfrak{S}^{\frac{1}{2}} \subset \mathfrak{S}^{-\frac{1}{2}}$ on $\mathfrak{S}$ where
$$\mathfrak{S}^{-\frac{1}{2}} := \mathfrak{S}\,, \qquad\qquad \mathfrak{S}^{\frac{1}{2}} = \langle \xi^A \rangle = \{\alpha^A \in \mathfrak{S} : \alpha^A \xi_A = 0\}\,.$$

The notion of principal spinors can defined in exactly the same way as in four dimensions: the spinor ξ^A defines a filtration
$$\mathfrak{S}^{\frac{k}{2}} \subset \mathfrak{S}^{\frac{k}{2}-1} \subset \ldots \subset \mathfrak{S}^{-\frac{k}{2}+1} \subset \mathfrak{S}^{-\frac{k}{2}}$$

---

[4] So, unlike in higher odd dimensions, the Lie algebra $\mathfrak{so}(3,\mathbb{C})$ admits a $|1|$-grading.

on $\mathfrak{S}^{-\frac{k}{2}} := \odot^k \mathfrak{S}$, where

$$\mathfrak{S}^{\frac{k-2\ell+2}{2}} := \left\{ \phi_{A_1 \ldots A_k} \in \mathfrak{S}^{-\frac{k}{2}} : \phi_{A_1 \ldots A_\ell A_{\ell+1} \ldots A_k} \xi^{A_1} \ldots \xi^{A_\ell} \right\} ,$$

and ξ^A is said to be a principal spinor of $\phi_{A_1 \ldots A_k}$ if it lies in $\mathfrak{S}^{-\frac{k}{2}+1}$.

In the particular case $k = 2$, we have a filtration $\mathfrak{g}^1 \subset \mathfrak{g}^0 \subset \mathfrak{g}^{-1}$ on $\mathfrak{g}$ the Lie algebra $\mathfrak{so}(3, \mathbb{C})$, or equivalently on $\mathfrak{V} \cong \wedge^2 \mathfrak{V} \cong \mathfrak{g}$, where

$$\mathfrak{g}^{-1} := \mathfrak{g} \,, \qquad \mathfrak{g}^0 := \left\{ \phi_{AB} \in \odot^2 \mathfrak{S} : \phi_{AB} \xi^A \xi^B = 0 \right\} \,, \qquad \mathfrak{g}^1 := \left\{ \phi_{AB} \in \odot^2 \mathfrak{S} : \phi_{AB} \xi^A = 0 \right\} \,.$$

Similarly, the case $k = 4$ yields filtrations on $\mathfrak{F}$ and $\mathfrak{A}$, but it is worth noting that unlike in higher dimensions, the filtration on $\mathfrak{A}$ is ‘shortened’ by two vector subspaces, i.e.

$$\{0\} =: \mathfrak{A}^3 \subset \mathfrak{A}^2 \subset \mathfrak{A}^1 \subset \mathfrak{A}^0 \subset \mathfrak{A}^{-1} \subset \mathfrak{A}^{-2} := \mathfrak{A} \,.$$

Foliating spinors and others The filtration on the $\mathfrak{p}$-module $\mathfrak{W} := \mathfrak{V} \otimes \mathfrak{g}/\mathfrak{p}$, where $\mathfrak{p} := \mathfrak{g}^0$ stabilises ξ^A, defining the classes of intrinsic torsion associated to the spinor field ξ^A is also shortened to $\mathfrak{W}^0 \subset \mathfrak{W}^{-1} \subset \mathfrak{W}^{-2}$. From a geometric point of view, this is reflected by the fact that on a 3-dimensional complex Riemannian manifold, the null plane $\mathcal{N}$ associated to a projective spinor field ξ^A is only 1-dimensional, and is thus trivially integrable. The remaining differential conditions then reduce to only three as given by the following proposition.

Proposition B.1 *Let $[\xi^A]$ be a holomorphic projective spinor field on a 3-dimensional complex Riemannian manifold with associated γ-plane distribution $\mathcal{N}$. Then the intrinsic torsion of $[\xi^A]$*

- *lies in $\mathfrak{W}^{-1}$ if and only if $\xi^A \xi^B \xi^C \nabla_{AB} \xi_C = 0$ if and only if $[\Gamma(\mathcal{N}^\perp), \Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N}^\perp)$;*
- *lies in $\mathfrak{W}^0$ if and only if $\xi^B \xi^C \nabla_{AB} \xi_C = 0$ if and only if $[\Gamma(\mathcal{N}^\perp), \Gamma(\mathcal{N}^\perp)] \subset \Gamma(\mathcal{N})$;*
- *vanishes if and only if $\xi^C \nabla_{AB} \xi_C = 0$.*

Remark B.2 The above conditions are equivalent to the null vector field $k^{AB} := \xi^A \xi^B$ being geodetic, dilation free and recurrent respectively.

B.2 Five dimensions

Let $(\mathfrak{V}, g)$ be a five-dimensional complex vector space equipped with a non-degenerate symmetric bilinear form. The double cover of the special orthogonal groups $\mathrm{SO}(5, \mathbb{C})$ is the complex symplectic group $\mathrm{Sp}(4, \mathbb{C})$. The spinor space $\mathfrak{S}$ and its dual space are thus four-dimensional complex equipped with non-degenerate symplectic forms ε_{AB} and ε^{AB} respectively. One can convert tensor indices into spinorial ones by means of the normalised skewsymmetric Van der Waerden γ-matrices $\frac{\mathrm{i}}{2} \gamma_a{}^{AB}$, tracefree with respect to ε_{AB}, which satisfy the identity

$$\gamma_{aA}{}^B \gamma^a{}_C{}^D = \delta_A^B \delta_C^D - 2 \delta_A^D \delta_C^B - 2 \varepsilon_{AC} \varepsilon^{BD} \,, \qquad \text{i.e.} \qquad \gamma_{aAB} \gamma^a{}_{CD} = \varepsilon_{AB} \varepsilon_{CD} + 4 \varepsilon_{A[C} \varepsilon_{D]B} \,. \tag{B.1}$$

In particular, we have $\mathfrak{V} \cong (\wedge^2 \mathfrak{S})_\circ$ and $\wedge^2 \mathfrak{V} \cong \odot^2 \mathfrak{S}$.

The tracefree Ricci tensor, the Weyl tensor and the Cotton tensor admit the spinorial expressions

$$\begin{aligned} \Phi_{ABCD} &= \Phi_{[AB][CD]} \,, & \Phi_{[ABC]D} &= 0 \,, \\ C_{ABCD} &= C_{(ABCD)} \,, & & \\ A_{ABCD} &= A_{[AB](CD)} \,, & A_{[ABC]D} &= 0 \,, \end{aligned}$$

respectively, all of which are completely tracefree.

Principal spinors Following the notation of section 2, a spinor ξ^A induces a filtration $\mathfrak{S}^1 \subset \mathfrak{S}^0 \subset \mathfrak{S}^{-1}$ on $\mathfrak{S}$ where

$$\mathfrak{S}^{-1} := \mathfrak{S}\,, \qquad \mathfrak{S}^0 := \{\alpha^A \in \mathfrak{S} : \alpha_A \xi^A = 0\}\,, \qquad \mathfrak{S}^1 := \langle \xi^A \rangle = \{\alpha^A \in \mathfrak{S} : \alpha_{[A}\xi_{B]} = 0\}\,,$$

In particular, the concept of a principal null spinor ξ^A of a spinorial object of any valence by means of the wedge and contraction operations.

The spinor ξ^A induces a filtration $\mathfrak{V}^1 \subset \mathfrak{V}^0 \subset \mathfrak{V}^{-1}$ where

$$\mathfrak{V}^{-1} := \mathfrak{V}\,, \qquad \mathfrak{V}^0 := \left\{ V^{AB} \in \mathfrak{V} : \xi_C V^{C[A}\,\xi^{B]} = 0 \right\}\,, \qquad \mathfrak{V}^1 := \left\{ V^{AB} \in \mathfrak{V} : \xi_C V^{CA} = 0 \right\}\,,$$

where[5] we can define equivalently $\mathfrak{V}^{-1} = \left\{ V^{AB} \in \mathfrak{V} : \xi_{[A} V_{BC]} = 0 \right\}$

Similarly, the irreducible graded structure of the Lie algebra $\mathfrak{so}(5,\mathbb{C})$ can be expressed in terms of kernels of the maps

$$\begin{aligned}
&{}^{\mathfrak{g}}\Pi^0_{-2}(\phi) := \xi^A \xi^B \phi_{AB}\,, && {}^{\mathfrak{g}}\Pi^0_{-1}(\phi) := \xi^A \phi_A{}^{[B}\,\xi^{C]}\,, && \\
&{}^{\mathfrak{g}}\Pi^0_0(\phi) := \xi^A \phi_A{}^B\,, && {}^{\mathfrak{g}}\Pi^1_0(\phi) := \xi_{[A}\,\phi_{B][C}\,\xi_{D]}\,, && {}^{\mathfrak{g}}\Pi^0_1(\phi) := \phi_{A[B}\,\xi_{C]}\,,
\end{aligned}$$

where $\phi_{AB} = \phi_{(AB)}$.

The explicit expressions for the maps ${}^{\mathfrak{F}}\Pi^j_i$, ${}^{\mathfrak{A}}\Pi^j_i$ and ${}^{\mathfrak{C}}\Pi^j_i$ defined in section 3 can be significantly simplified. For $\Phi_{ABCD} \in \mathfrak{F}$, we have

$$\begin{aligned}
&{}^{\mathfrak{F}}\Pi^0_{-2}(\Phi) := \xi_{[F}\,\Phi_{A]BC[D}\,\xi^B \xi^C \xi_{E]}\,, && {}^{\mathfrak{F}}\Pi^0_{-1}(\Phi) := \Phi_{ABC[D}\,\xi^B \xi^C \xi_{E]}\,, \\
&{}^{\mathfrak{F}}\Pi^0_0(\Phi) := \Phi_{ABCD} \xi^B \xi^C\,, && {}^{\mathfrak{F}}\Pi^1_0(\Phi) := \xi_{[A}\Phi_{B]ECD}\xi^E + \varepsilon_{[A|[C}\Phi_{D]EF|B]}\xi^E \xi^F\,, \\
&{}^{\mathfrak{F}}\Pi^0_1(\Phi) := \Phi_{ABCD}\xi^B\,. &&
\end{aligned}$$

For $A_{ABCD} \in \mathfrak{A}$, we have

$$\begin{aligned}
&{}^{\mathfrak{A}}\Pi^0_{-3}(A) := \xi_{[E}\,A_{A]BCD}\xi^B \xi^C \xi^D\,, && \\
&{}^{\mathfrak{A}}\Pi^0_{-2}(A) := A_{ABCD}\xi^B \xi^C \xi^D\,, && {}^{\mathfrak{A}}\Pi^1_{-2}(A) := \xi_{[A}A_{B]EF[C}\xi^E \xi^F \xi_{D]} + \xi_{[C}A_{D]EF[A}\xi^E \xi^F \xi_{B]}\,, \\
&{}^{\mathfrak{A}}\Pi^0_{-1}(A) := 4\xi_{[A}A_{B]DEC}\xi^D \xi^E - A_{ABDE}\xi^D \xi^E \xi_C\,, && {}^{\mathfrak{A}}\Pi^1_{-1}(A) := A_{ABC[D}\,\xi^B \xi^C \xi_{E]}\,, \\
&{}^{\mathfrak{A}}\Pi^3_{-1}(A) := \xi_{[A}\,A_{B]G[C\,|[E}\,\xi^G \xi_{F]}\xi_{|\,D]} + \xi_{[C}\,A_{D]G[E\,|[A}\,\xi^G \xi_{B]}\xi_{|\,F]} + \xi_{[E}\,A_{F]G[A\,|[C}\,\xi^G \xi_{D]}\xi_{|\,B]}\,, && \\
&{}^{\mathfrak{A}}\Pi^0_0(A) := A_{ABCD}\xi^B \xi^C\,, && {}^{\mathfrak{A}}\Pi^1_0(A) := \xi^F A_{AF[B|[D}\xi_{E]}\xi_{|C]}\,, \\
&{}^{\mathfrak{A}}\Pi^2_0(A) := \xi_{[A}\xi^F A_{B]FE[C}\xi_{D]} + \xi_{[A}\xi^F A_{B]EF[C}\xi_{D]}\,, && \\
&{}^{\mathfrak{A}}\Pi^0_1(A) := \xi_{[A}\xi^E A_{B]ECD} - \xi^E A_{ABE(C}\xi_{D)}\,, && {}^{\mathfrak{A}}\Pi^1_1(A) := \xi^E A_{AEB[C}\xi_{D]} \\
&{}^{\mathfrak{A}}\Pi^3_1(A) := A_{AB[C\,|[E}\,\xi_{F]}\xi_{|\,D]} + A_{CD[E\,|[A}\,\xi_{B]}\xi_{|\,F]} + A_{EF[A\,|[C}\,\xi_{D]}\xi_{|\,B]}\,, && \\
&{}^{\mathfrak{A}}\Pi^0_2(A) := \xi^D A_{ADBC}\,, && {}^{\mathfrak{A}}\Pi^1_2(A) := A_{ABC[D}\xi_{E]} + A_{DEC[A}\xi_{B]}\,.
\end{aligned}$$

---

[5] This can be seen by means of a volume form ε_{ABCD} so that $\varepsilon_{ABCD} = 3\varepsilon_{[A\,B}\varepsilon_{C\,D]} = \frac{3}{5}\gamma_{a[A\,B}\gamma^a{}_{C\,D]}$, which can be seen to satisfy the normalisation so that $\varepsilon_{ABCD}\varepsilon^{EFGH} = 24\delta^E_{[A}\delta^F_B\delta^G_C\delta^H_{D]}$. Also, a vector (or 1-form) V^a is null if and only if it is of the form $V^{AB} = \xi^{[A}\,\alpha^{B]}$ for some spinors ξ_A and α_A with $\alpha_A\xi^A = 0$.

Finally, for $C_{ABCD} \in \mathfrak{C}$, we have

$$
\begin{aligned}
&{}^{\mathfrak{C}}\Pi^0_{-4}(C) := C_{ABCD}\xi^A\xi^B\xi^C\xi^D\,, && {}^{\mathfrak{C}}\Pi^0_{-3}(C) := C_{ABC[D}\,\xi^A\xi^B\xi^C\xi_{E]}\,, && \\
&{}^{\mathfrak{C}}\Pi^0_{-2}(C) := C_{ABCD}\xi^A\xi^B\xi^C\,, && {}^{\mathfrak{C}}\Pi^2_{-2}(C) := \xi_{[F}\,C_{A]BC[D}\,\xi^B\xi^C\xi_{E]}\,, && \\
&{}^{\mathfrak{C}}\Pi^0_{-1}(C) := C_{ABC[D}\,\xi^B\xi^C\xi_{E]}\,, && {}^{\mathfrak{C}}\Pi^1_{-1}(C) := \xi_{[F}\,C_{A]B[C\,|[D}\,\xi^B\xi_{E]}\xi_{|\,F]}\,, && \\
&{}^{\mathfrak{C}}\Pi^0_{0}(C) := C_{ABCD}\xi^B\xi^C\,, && {}^{\mathfrak{C}}\Pi^1_{0}(C) := \xi_{[F}\,C_{A]BC[D}\,\xi^B\xi_{F]}\,, && {}^{\mathfrak{C}}\Pi^4_{0}(C) := \xi_{[G}\,\xi_{|[F}\,C_{A]|\,B][C\,|[D}\,\xi_{E]}\xi_{|\,F]}\,, \\
&{}^{\mathfrak{C}}\Pi^0_{1}(C) := \xi_{[F}\,C_{A]BCD}\xi^B\,, && {}^{\mathfrak{C}}\Pi^1_{1}(C) := \xi_{[F}\,C_{A]B[C\,|[D}\,\xi_{E]}\xi_{|\,F]}\,, && \\
&{}^{\mathfrak{C}}\Pi^0_{2}(C) := C_{ABCD}\xi^B\,, && {}^{\mathfrak{C}}\Pi^2_{2}(C) := \xi_{[F}\,C_{A]BC[D}\,\xi_{E]}\,, && \\
&{}^{\mathfrak{C}}\Pi^0_{3}(C) := C_{ABC[D}\,\xi_{E]}\,. && &&
\end{aligned}
$$

B.2.1 Foliating and not so foliating spinors

On a five-dimensional complex Riemannian manifold $(\mathcal{M}, g)$, the differential characterisations (5.6), (5.7), (5.8), (5.9), (5.15), (5.12), (5.13), (5.14) of a projective spinor field ξ^A can be expressed as

$$\xi_{[A}\,\left(\xi^C\nabla_{B]C}\xi^D\right)\xi_D = 0\,, \tag{B.2}$$

$$\left(\xi^B\nabla_{AB}\xi^C\right)\xi_C = 0\,, \tag{B.3}$$

$$\left(\xi^D\nabla_{D[A}\xi_B\right)\xi_{C]} = 0\,, \tag{B.4}$$

$$\xi_{[A}\,\left(\xi^E\nabla_{B]E}\xi_{[C}\right)\xi_{D]} + \xi_{[C}\,\left(\xi^E\nabla_{D]E}\xi_{[A}\right)\xi_{B]} = 0\,, \tag{B.5}$$

$$(\nabla_{AB}\xi^C)\xi_C = 0\,, \tag{B.6}$$

$$\left(\xi^D\nabla_{AD}\xi_{[B}\right)\xi_{C]} = 0\,, \tag{B.7}$$

$$(\nabla_{AC}\xi^C)\xi^B - (\xi^C\nabla_{AC}\xi^B) = 0\,, \tag{B.8}$$

$$\left(\nabla_{AB}\xi_{[C}\right)\xi_{D]} + \xi_{[C}\varepsilon_{D][A}\nabla_{B]E}\xi^E + \varepsilon_{[C|[A}\xi^E\nabla_{B]E}\xi_{|D]} = 0\,, \tag{B.9}$$

respectively.

Further, condition (5.16) for an almost foliating spinor reduces to (B.2). Condition (5.17) for a $\gamma^\perp$-foliating spinor can be expressed as

$$\xi_{[A}\left(\xi^E\nabla_{B]E}\xi_{[C}\right)\xi_{D]} + \xi_{[C}\varepsilon_{D][A}\left(\xi^E\nabla_{B]E}\xi_F\right)\xi^F = 0\,. \tag{B.10}$$

while condition (5.18) for a foliating spinor

$$\left(\xi^B\nabla_{AB}\xi^C\right)\xi_C = 0\,, \qquad\qquad \xi_{[A}\,\left(\xi^E\nabla_{B]E}\xi_{[C}\right)\xi_{D]} = 0\,. \tag{B.11}$$

As an example, one can check that a solution ξ^A of the twistor equation

$$\nabla_{AB}\xi_C + \frac{1}{5}\varepsilon_{AB}\zeta_C + \frac{4}{5}\zeta_{[A}\,\varepsilon_{B]C} = 0\,, \qquad\qquad \zeta^A = \nabla^{AB}\xi_B\,,$$

satisfies equations (B.11) if and only if $\xi^A\zeta_A = 0$ as claimed in Proposition 5.21.

C Conformal structures

In this appendix, we collect a few facts, notations, and conventions pertaining to conformal and projective geometries. We roughly follow [BEG94], although our staggering of indices differs from theirs.

A *smooth, respectively holomorphic, conformal structure* on a smooth, respectively complex, manifold $\mathcal{M}$ is an equivalence class of smooth, respectively holomorphic, metrics $[g_{ab}]$ on $\mathcal{M}$, whereby two metrics $\hat{g}_{ab}$ and g_{ab} belong to the same class if and only if

$$\hat{g}_{ab} = \Omega^2 g_{ab}\,, \tag{C.1}$$

for some non-vanishing smooth, respecitvely homomorphic, function Ω on $\mathcal{M}$. The respective Levi-Civita connections ∇_a and $\hat{\nabla}_a$ of g_{ab} and $\hat{g}_{ab}$ are then related by

$$\hat{\nabla}_a V^b = \nabla_a V^b + Q_{ac}{}^b V^c \,, \qquad\qquad Q_{abc} := Q_{ab}{}^d g_{dc} = 2 \Upsilon_{(a} g_{b)c} - \Upsilon_c g_{ab} \,,$$

where $\Upsilon_a := \Omega^{-1} \nabla_a \Omega$.

Spinor bundles For a formal definition of the spinor bundle $\mathfrak{S}$ on a conformal manifold, see e.g. [BJ10]. For our purpose, we simply note that under a rescaling (C.1), the Van der Waerden γ-matrices can be chosen to transform as

$$\gamma_{aA}{}^B \mapsto \hat{\gamma}_{aA}{}^B = \Omega \gamma_{aA}{}^B \,,$$

where $\hat{\gamma}_{aA}{}^B$ denote the Van der Waerden γ-matrices for metric $\hat{g}_{ab}$. In addition, we can choose the spin invariant bilinear forms ε_{AB} on $\mathcal{S}$ to rescale with a conformal weight of 1, and their dual with a conformal weight of -1. This means in particular that the quantities $\gamma_a{}^{AB}$ and $\gamma^a{}_{AB}$ have conformal weight 0. Then the spin connection $\hat{\nabla}_a$ is related to ∇_a by

$$\begin{aligned} \hat{\nabla}_a \xi^B &= \nabla_a \xi^B - \frac{1}{2} \Upsilon_b \gamma^b{}_{aC}{}^B \xi^C + \frac{1}{2} \Upsilon_a \xi^B \\ &= \nabla_a \xi^B - \frac{1}{2} \Upsilon_b \gamma^b{}_C{}^D \gamma_{aD}{}^B \xi^C \,, \end{aligned} \tag{C.2}$$

and similarly for dual spinors. This connection preserves the hatted γ-matrices *and* the hatted bilinear forms on $\mathcal{S}$, in agreement with the convention of [PR84].

If we now assume that ξ^A is a pure spinor field, then we obtain from (C.2)

$$\begin{aligned} \left(\hat{\nabla}_a \hat{\xi}^{bB} \right) \hat{\xi}_b{}^C &= \left(\nabla_a \xi^{bB} \right) \xi_b{}^C + \frac{1}{2} \Upsilon_b \xi^{bD} \gamma_{aD}{}^B \xi^C + 2 \Upsilon_b \xi^{b[B} \xi_a{}^{C]} \,, \\ \left(\hat{\xi}^{aA} \hat{\nabla}_a \xi^B \right) &= \Omega^{-1} \left(\xi^{aA} \nabla_a \xi^B - \frac{1}{2} \Upsilon_b \xi^{bB} \xi^A + \Upsilon_b \xi^{bA} \xi^B \right) \,, \\ \left(\hat{\xi}^{aA} \hat{\nabla}_a \hat{\xi}^{bB} \right) \hat{\xi}_b{}^C &= \Omega^{-1} \left(\left(\xi^{aA} \nabla_a \xi^{bB} \right) \xi_b{}^C - \frac{1}{2} \Upsilon_b \xi^{bB} \xi^A \xi^C - 2 \Upsilon_b \xi^{b[A} \xi^{C]} \xi^B \right) \,, \\ \left(\hat{\nabla}_a \hat{\xi}^{aB} \right) \xi^C - \hat{\xi}^{aB} \hat{\nabla}_a \xi^C &= \Omega^{-1} \left(\left(\nabla_a \xi^{aB} \right) \xi^C - \xi^{aB} \nabla_a \xi^C + \frac{n-2}{2} \Upsilon_b \xi^{bB} \xi^C + \frac{1}{2} \Upsilon_b \xi^{bC} \xi^B \right) \,, \end{aligned}$$

where we have set $\hat{\xi}^{aA} := \hat{\gamma}^a{}_B{}^A \xi^B$. Combining the first and last of these expressions yields

$$\begin{aligned} &\left(\hat{\nabla}_a \hat{\xi}^{b[B} \right) \hat{\xi}_b{}^C \xi^{D]} - \frac{1}{2(m-1)} \left(\left(\hat{\nabla}_b \hat{\xi}^{b[B} \right) \xi^C - \hat{\xi}^{b[B} \hat{\nabla}_b \xi^C \right) \hat{\xi}_a{}^{D]} \\ &\qquad\qquad = \left(\nabla_a \xi^{b[B} \right) \xi_b{}^C \xi^{D]} - \frac{1}{2(m-1)} \left(\left(\nabla_b \xi^{b[B} \right) \xi^C - \xi^{b[B} \nabla_b \xi^C \right) \xi_a{}^{D]} \,. \end{aligned}$$

Curvature The conformal properties of the Riemann curvature can be conveniently brought out by means of the decomposition

$$R_{abcd} = C_{abcd} - 4 g_{[c\,|[a} \mathrm{P}_{b]|\,d]} \,, \qquad\qquad \mathrm{P}_{ab} := \frac{1}{2-n} \Phi_{ab} - R \frac{1}{2n(n-1)} g_{ab} \,. \tag{C.3}$$

where the Weyl tensor C_{abcd} is the conformally invariant part of R_{abcd}, and the *Schouten* or *Rho tensor* P_{ab}, together with its its trace P, transform as

$$\hat{\mathrm{P}}_{ab} = \mathrm{P}_{ab} - \nabla_a \Upsilon_b + \Upsilon_a \Upsilon_b - \frac{1}{2} \Upsilon_c \Upsilon^c g_{ab} \,, \qquad \hat{\mathrm{P}} = \Omega^{-2} \left(\mathrm{P} - \nabla^c \Upsilon_c - \frac{n-2}{2} \Upsilon^c \Upsilon_c \right) \,. \tag{C.4}$$

Finally, the Cotton-York tensor

$$A_{abc} := 2\nabla_{[b}\mathrm{P}_{c]a} = -(n-3)\nabla^d C_{dabc} \,,$$

where the expression on the RHS follows from the contracted Bianchi identity, transforms as

$$\hat{A}_{abc} = A_{abc} - \Upsilon^d C_{dabc} \,.$$

D Tables of irreducibles

In this appendix, we have collected tables of the irreducible $\mathfrak{g}$-modules and their $\mathfrak{p}$-modules decomposition in terms of their dominant weights and dimensions. Much of this representation theoretic description is already explained in [TC12b], and more thoroughly in [BE89, ČS09b]. We shall therefore remain concise on this subject, and refer the reader to the aforementioned references for more details.

D.1 The Lie algebra $\mathfrak{so}(2m+1,\mathbb{C})$

The Dynkin diagram for $\mathfrak{g} := \mathfrak{so}(2m+1,\mathbb{C})$ is

 , m nodes.

In the case $m=2$, this coincides with the 'reversed' Dynkin diagram  of $\mathfrak{sp}(4,\mathbb{C})$.

The irreducible $\mathfrak{g}$-modules considered in this article can be represented in terms of their dominant weights and dimensions as given in the table below:

$\mathfrak{g}$-mod	Dominant weight	Dimension
$\mathfrak{V}$		$2m+1$
$\mathfrak{g}$		$m(2m+1)$
$\mathfrak{F}$		$m(2m+3)$
$\mathfrak{A}$		$\frac{1}{3}(2m-1)(2m+1)(2m+3)$
$\mathfrak{C}$		$\frac{1}{3}(m-1)(m+1)(2m+1)(2m+3)$

D.2 The parabolic Lie subalgebra $\mathfrak{p}$

To the (standard) parabolic Lie subalgebra $\mathfrak{p}$ stabilising a pure spinor ξ^A corresponds the mutilated Dynkin diagram

 .

The Dynkin diagram for the reductive part $\mathfrak{g}_0 \cong \mathfrak{gl}(m,\mathbb{C})$ can be recovered by eliminating the crossed node.

We have collected tables of the irreducibles $\mathfrak{p}$-modules occuring in the $\mathfrak{p}$-invariant decompositions of the representations $\mathfrak{V}$, $\mathfrak{g}$, $\mathfrak{F}$, $\mathfrak{A}$, $\mathfrak{C}$ and $\mathfrak{W}$. Care must be taken in five dimensions where the modules $\mathfrak{A}^2_{\pm 1}$, $\mathfrak{C}^2_{\pm 2}$, $\mathfrak{C}^1_{\pm 1}$, $\mathfrak{C}^3_0$ do not occur, and the isotopic modules $\mathfrak{C}^1_0$ and $\mathfrak{C}^2_0$ degenerate to the same module. In dimension seven, the module $\mathfrak{C}^3_0$ does not occur.

Acknowlegments This work is funded by a SoMoPro (South Moravian Programme) Fellowship: it has received a financial contribution from the European Union within the Seventh Framework Programme (FP/2007-2013) under Grant Agreement No. 229603, and is also co-financed by the South Moravian Region.

$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dominant weight	$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dominant weight	Dimension
$\mathfrak{V}_1$	$\mathfrak{V}_1$	$\overset{1}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\Rightarrow\overset{0}{\times}$	$\mathfrak{V}^{-1}/\mathfrak{V}^0$	$\mathfrak{V}_{-1}$	$\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{1}{\bullet}\Rightarrow\overset{-2}{\times}$	m
$\mathfrak{V}^0/\mathfrak{V}^1$	$\mathfrak{V}_0$	$\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\Rightarrow\overset{0}{\times}$				1

Table 1: Irreducible $\mathfrak{p}$-modules of $\mathfrak{V}$

$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dominant weight	$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dominant weight	Dimension
$\mathfrak{g}^0_2$	$\mathfrak{g}_2$	$\overset{0}{\bullet}\!-\!\overset{1}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\Rightarrow\overset{0}{\times}$	$\mathfrak{g}^0_{-2}$	$\mathfrak{g}_{-2}$	$\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{1}{\bullet}\!-\!\overset{0}{\bullet}\Rightarrow\overset{-2}{\times}$	$\frac{1}{2}m(m-1)$
$\mathfrak{g}^0_1$	$\mathfrak{g}_1$	$\overset{1}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\Rightarrow\overset{0}{\times}$	$\mathfrak{g}^0_{-1}$	$\mathfrak{g}_{-1}$	$\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{1}{\bullet}\Rightarrow\overset{-2}{\times}$	m
$\mathfrak{g}^0_0$	$\mathfrak{z}_0$	$\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\Rightarrow\overset{0}{\times}$				1
$\mathfrak{g}^1_0$	$\mathfrak{sl}_0$	$\overset{1}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\cdots\overset{0}{\bullet}\!-\!\overset{0}{\bullet}\!-\!\overset{1}{\bullet}\Rightarrow\overset{-2}{\times}$				m^2-1

Table 2: Irreducible $\mathfrak{p}$-modules of $\mathfrak{V}$

References

[AG97] V. Apostolov and P. Gauduchon, *The Riemannian Goldberg-Sachs theorem*, Internat. J. Math. **8** (1997), no. 4, 421–439.

[BE89] Robert J. Baston and Michael G. Eastwood, *The Penrose transform*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1989. Its interaction with representation theory, Oxford Science Publications.

[BEG94] T. N. Bailey, M. G. Eastwood, and A. R. Gover, *Thomas's structure bundle for conformal, projective and related structures*, Rocky Mountain J. Math. **24** (1994), no. 4, 1191–1217.

[BJ10] Helga Baum and Andreas Juhl, *Conformal differential geometry*, Oberwolfach Seminars, vol. 40, Birkhäuser Verlag, Basel, 2010. Q-curvature and conformal holonomy. MR2598414 (2011d:53055)

[Bry00] Robert L. Bryant, *Pseudo-Riemannian metrics with parallel spinor fields and vanishing Ricci tensor*, Global analysis and harmonic analysis (Marseille-Luminy, 1999), 2000, pp. 53–94. MR1822355 (2002h:53082)

[BT89] Paolo Budinich and Andrzej Trautman, *Fock space description of simple spinors*, J. Math. Phys. **30** (1989), no. 9, 2125–2131.

[Car10] Elie Cartan, *Les systèmes de Pfaff, à cinq variables et les équations aux dérivées partielles du second ordre*, Ann. Sci. École Norm. Sup. (3) **27** (1910), 109–192. MR1509120

[Car81] Élie Cartan, *The theory of spinors*, Dover Publications Inc., New York, 1981. With a foreword by Raymond Streater, A reprint of the 1966 English translation, Dover Books on Advanced Mathematics. MR631850 (83a:15017)

[CMPP04] A. Coley, R. Milson, V. Pravda, and A. Pravdová, *Classification of the Weyl tensor in higher dimensions*, Classical Quantum Gravity **21** (2004), no. 7, L35–L41, available at `arXiv:gr-qc/0401008`.

[ČS09a] Andreas Čap and Katja Sagerschnig, *On Nurowski's conformal structure associated to a generic rank two distribution in dimension five*, J. Geom. Phys. **59** (2009), no. 7, 901–912. MR2536853 (2010f:58003)

[ČS09b] Andreas Čap and Jan Slovák, *Parabolic geometries. I*, Mathematical Surveys and Monographs, vol. 154, American Mathematical Society, Providence, RI, 2009. Background and general theory. MR2532439 (2010j:53037)

[DS05] Pieter-Jan De Smet, *The Petrov type of the BMPV metric*, Gen. Rel. Grav. **37** (2005), 237–242, available at `arXiv:gr-qc/0401033`.

[Eas95] Michael G. Eastwood, *A twistor description of null self-dual maxwell fields*, Further advances in twistor theory. Vol. II, 1995, pp. 96–99.

$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dimension
$\mathfrak{F}^0_{\pm2}$	$\mathfrak{V}_{\pm1} \circledcirc \mathfrak{V}_{\pm1}$	$\frac{1}{2}m(m+1)$
$\mathfrak{F}^0_{\pm1}$	$\mathfrak{V}_0 \circledcirc \mathfrak{V}_{\pm1}$	m
$\mathfrak{F}^0_0$	$\mathfrak{V}_{\pm1} \circledcirc \mathfrak{V}_{\mp1}$	1
$\mathfrak{F}^1_0$	$\mathfrak{V}_0 \circledcirc \mathfrak{V}_0$	$m(m-1)$

Table 3: Irreducible $\mathfrak{p}$-modules of $\mathfrak{F}$

$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dimension
$\mathfrak{A}^0_{\pm3}$	$V_{\pm1} \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{3}m(m+1)(m-1)$
$\mathfrak{A}^0_{\pm2}$	$V_0 \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{2}m(m-1)$
$\mathfrak{A}^1_{\pm2}$	$V_{\pm1} \circledcirc \mathfrak{g}_{\pm1}$	$\frac{1}{2}m(m+1)$
$\mathfrak{A}^0_{\pm1}$	$V_{\pm1} \circledcirc \mathfrak{z}_0$	m
$\mathfrak{A}^1_{\pm1}$	$V_0 \circledcirc \mathfrak{g}_{\pm1}$	m
$\mathfrak{A}^2_{\pm1}$	$V_{\mp1} \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{2}m(m-2)(m+1)$
$\mathfrak{A}^3_{\pm1}$	$V_{\pm1} \circledcirc \mathfrak{sl}_0$	$\frac{1}{2}m(m+2)(m-1)$
$\mathfrak{A}^0_0$	$V_0 \circledcirc \mathfrak{z}_0$	1
$\mathfrak{A}^1_0$	$V_0 \circledcirc \mathfrak{sl}_0$	$(m-1)(m+1)$
$\mathfrak{A}^2_0$	$V_1 \circledcirc \mathfrak{g}_{-1}$	$(m-1)(m+1)$

Table 4: Irreducible $\mathfrak{p}$-modules of $\mathfrak{A}$

$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dimension
$\mathfrak{C}^0_{\pm4}$	$\mathfrak{g}_{\pm2} \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{12}m^2(m^2-1)$
$\mathfrak{C}^0_{\pm3}$	$\mathfrak{g}_{\pm1} \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{3}m(m^2-1)$
$\mathfrak{C}^0_{\pm2}$	$\mathfrak{z}_0 \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{2}m(m-1)$
$\mathfrak{C}^1_{\pm2}$	$\mathfrak{g}_{\pm1} \circledcirc \mathfrak{g}_{\pm1}$	$\frac{1}{2}m(m+1)$
$\mathfrak{C}^2_{\pm2}$	$\mathfrak{sl}_0 \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{3}m^2(m^2-4)$
$\mathfrak{C}^0_{\pm1}$	$\mathfrak{z}_0 \circledcirc \mathfrak{g}_{\pm1}$	m
$\mathfrak{C}^1_{\pm1}$	$\mathfrak{g}_{\mp1} \circledcirc \mathfrak{g}_{\pm2}$	$\frac{1}{2}m(m-2)(m+1)$
$\mathfrak{C}^2_{\pm1}$	$\mathfrak{sl}_0 \circledcirc \mathfrak{g}_{\pm1}$	$\frac{1}{2}m(m+2)(m-1)$
$\mathfrak{C}^0_0$	$\mathfrak{z}_0 \circledcirc \mathfrak{z}_0$	1
$\mathfrak{C}^1_0$	$\mathfrak{sl}_0 \circledcirc \mathfrak{z}_0$	m^2-1
$\mathfrak{C}^2_0$	$\mathfrak{g}_1 \circledcirc \mathfrak{g}_{-1}$	m^2-1
$\mathfrak{C}^3_0$	$\mathfrak{g}_2 \circledcirc \mathfrak{g}_{-2}$	$\frac{1}{4}m^2(m+1)(m-3)$
$\mathfrak{C}^4_0$	$\mathfrak{sl}_0 \circledcirc \mathfrak{sl}_0$	$\frac{1}{4}m^2(m-1)(m+3)$

Table 5: Irreducible $\mathfrak{p}$-modules of $\mathfrak{C}$.

[FFS94] M. Falcitelli, A. Farinola, and S. Salamon, *Almost-Hermitian geometry*, Differential Geom. Appl. **4** (1994), no. 3, 259–282. MR1299398 (95i:53047)

[FH91] William Fulton and Joe Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991. A first course, Readings in Mathematics.

[GH80] Alfred Gray and Luis M. Hervella, *The sixteen classes of almost Hermitian manifolds and their linear invariants*, Ann. Mat. Pura Appl. (4) **123** (1980), 35–58. MR581924 (81m:53045)

[GHN10] A. Gover, C. Hill, and Paweł Nurowski, *Sharp version of the Goldberg-Sachs theorem*, Annali di Matematica Pura ed Applicata (2010), 1–46. 10.1007/s10231-010-0151-4.

[God10] Mahdi Godazgar, *Spinor classification of the Weyl tensor in five dimensions*, Classical Quantum Gravity **27** (2010), no. 24, 245013, 37. MR2739969

[GPGLMG09] Alfonso García-Parrado Gómez-Lobo and José M. Martín-García, *Spinor calculus on five-dimensional spacetimes*, J. Math. Phys. **50** (2009), no. 12, 122504, 26. MR2582586

[GS09] J. Goldberg and R. Sachs, *Republication of: A theorem on Petrov types*, General Relativity and Gravitation **41** (2009), 433–444. 10.1007/s10714-008-0722-5.

[HM88] Lane P. Hughston and Lionel J. Mason, *A generalised Kerr-Robinson theorem*, Classical Quantum Gravity **5** (1988), no. 2, 275–285.

[HS09] Matthias Hammerl and Katja Sagerschnig, *Conformal structures associated to generic rank 2 distributions on 5-manifolds—characterization and Killing-field decomposition*, SIGMA Symmetry Integrability Geom. Methods Appl. **5** (2009), Paper 081, 29. MR2529166 (2011b:58009)

$\mathfrak{p}$-mod	$\mathfrak{g}_0$-mod	Dominant weight	Dimension
$\mathfrak{W}^0_{-3}$	$\wedge^3\mathfrak{V}_{-1}$		$\frac{1}{3!}m(m-1)(m-2)$
$\mathfrak{W}^1_{-3}$	$\mathfrak{V}_{-1} \circledcirc \mathfrak{g}_{-2}$		$\frac{1}{3}m(m^2-1)$
$\mathfrak{W}^0_{-2}$	$\mathfrak{V}_0 \otimes \mathfrak{g}_{-2}$		$\frac{1}{2}m(m-1)$
$\mathfrak{W}^1_{-2}$	$\wedge^2\mathfrak{V}_{-1} \otimes \mathfrak{V}_0$		$\frac{1}{2}m(m-1)$
$\mathfrak{W}^2_{-2}$	$\odot^2\mathfrak{V}_{-1} \otimes \mathfrak{V}_0$		$\frac{1}{2}m(m+1)$
$\mathfrak{W}^0_{-1}$	$\mathfrak{z}_0 \otimes \mathfrak{V}_{-1}$		m
$\mathfrak{W}^1_{-1}$	$\mathfrak{sl}_0 \otimes_\circ \mathfrak{V}_{-1}$		$\frac{1}{2}m(m+1)(m-2)$
$\mathfrak{W}^2_{-1}$	$\mathfrak{V}_0 \otimes \mathfrak{g}_{-1}$		m
$\mathfrak{W}^0_0$	$\mathfrak{z}_0 \otimes \mathfrak{V}_0$		1
$\mathfrak{W}^1_0$	$\mathfrak{sl}_0 \otimes \mathfrak{V}_0$		m^2-1

Table 6: Irreducible $\mathfrak{p}$-modules of $\mathfrak{W}$

[HS11] ______, *The twistor spinors of generic 2- and 3-distributions*, Ann. Global Anal. Geom. **39** (2011), no. 4, 403–425. MR2776770

[HS95] John P. Harnad and S. Shnider, *Isotropic geometry and twistors in higher dimensions. 2: Odd dimensions, reality conditions, and twistor superspaces*, J. Math. Phys. **36** (1995), 1945–1970.

[Hug95] Lane P. Hughston, *Differential geometry in six dimensions*, Further advances in twistor theory. Vol. II, 1995, pp. 79–82.

[Jef95] B. P. Jeffryes, *A six-dimensional 'Penrose diagram'*, Further advances in twistor theory. Vol. II, 1995, pp. 85–87.

[Kop97] Wojciech Kopczyński, *Pure spinors in odd dimensions*, Classical Quantum Gravity **14** (1997), no. 1A, A227–A236. Geometry and physics.

[LM89] H. B. Lawson Jr. and M.-L. Michelsohn, *Spin geometry*, Princeton Mathematical Series, vol. 38, Princeton University Press, Princeton, NJ, 1989.

[MCPP05] R. Milson, A. Coley, V. Pravda, and A. Pravdová, *Alignment and algebraically special tensors in Lorentzian geometry*, Int. J. Geom. Methods Mod. Phys. **2** (2005), no. 1, 41–61, available at `arXiv:gr-qc/0401010`.

[MT10] L. Mason and A. Taghavi-Chabert, *Killing-Yano tensors and multi-Hermitian structures*, Journal of Geometry and Physics **60** (June 2010), 907–923, available at `arXiv:0805.3756`.

[NT02] P. Nurowski and A. Trautman, *Robinson manifolds as the Lorentzian analogs of Hermite manifolds*, Differential Geom. Appl. **17** (2002), no. 2-3, 175–195. 8th International Conference on Differential Geometry and its Applications (Opava, 2001).

[Nur05] Paweł Nurowski, *Differential equations and conformal structures*, J. Geom. Phys. **55** (2005), no. 1, 19–49. MR2157414 (2006k:53021)

[Pen60] Roger Penrose, *A spinor approach to general relativity*, Ann. Physics **10** (1960), 171–201. MR0115765 (22 #6563)

[Pet00] A. Z. Petrov, *The classification of spaces defining gravitational fields*, Gen. Relativity Gravitation **32** (2000), no. 8, 1665–1685. Reprint of Kazan. Gos. Univ. Uč. Zap. **1**14 (1954), no. 8, 55–69 [MR0076401 (17,892g)]. MR1784371 (2002h:83028)

[PR84] Roger Penrose and Wolfgang Rindler, *Spinors and space-time. Vol. 1*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1984. Two-spinor calculus and relativistic fields.

[PR86] ______, *Spinors and space-time. Vol. 2*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1986. Spinor and twistor methods in space-time geometry.

[Rob61] Ivor Robinson, *Null electromagnetic fields*, J. Mathematical Phys. **2** (1961), 290–291.

[Sal89] Simon Salamon, *Riemannian geometry and holonomy groups*, Pitman Research Notes in Mathematics Series, vol. 201, Longman Scientific & Technical, Harlow, 1989. MR1004008 (90g:53058)

[TC11] Arman Taghavi-Chabert, *Optical structures, algebraically special spacetimes, and the Goldberg-Sachs theorem in five dimensions*, Class. Quant. Grav. **28** (2011), 145010 (32pp), available at `1011.6168`.

[TC12a] ———, *The complex Goldberg-Sachs theorem in higher dimensions*, J. Geom. Phys. **62** (2012), no. 5, 981–1012. MR2901842

[TC12b] ———, *Pure spinors, intrinsic torsion and curvature in even dimensions* (2012), available at `1212.3595`.

[TC] ———, *The curvature of almost Robinson manifolds*. In preparation.

[Tra02] Andrzej Trautman, *Robinson manifolds and Cauchy-Riemann spaces*, Classical Quantum Gravity **19** (2002), no. 2, R1–R10.

[TV81] Franco Tricerri and Lieven Vanhecke, *Curvature tensors on almost Hermitian manifolds*, Trans. Amer. Math. Soc. **267** (1981), no. 2, 365–397. MR626479 (82j:53071)

[Wit59] Louis Witten, *Invariants of general relativity and the classification of spaces*, Phys. Rev. (2) **113** (1959), 357–362. MR0115764 (22 #6562)